

New models for sustainable agriculture





www.barillacfn.com info@barillacfn.com

Advisory Board Barbara Buchner, Claude Fischler, Mario Monti, John Reilly Gabriele Riccardi, Camillo Ricordi, Umberto Veronesi

In collaboration with The European House-Ambrosetti

Publishing and editorial coordination Codice Edizioni

Art direction and layout adfarmandchicas

New models for sustainable agriculture (November 2011)

Images National Geographic Image Collection Corbis Images

Cover image: Corbis





ear Reader,
We live in an era when agriculture is demonstrating its fragility on a daily basis.

Despite its high complexity, for decades we have been accustomed to regard it as a field of activities that deliver modest added value and poor technical, avant-garde content and free from risk of structural discontinuity. As its importance has gradually diminished within the economies of Western countries, the degree of public attention has also declined.

The awakening could not have been more abrupt: for some years now, market crises, the decrease in the productivity growth rates and the emergence of serious distribution problems have brought the agri-food sector to the forefront of international political and economic debate.

In particular, concerns are emerging around the level of exposure to possible structural shocks of the global food chain. It is increasingly clear that agriculture will have to search for and find a new medium-long term balance in order to withstand the great phenomena of demographic, climate, geopolitical and economic change affecting the world.

Our interest in the subject of agricultural models stems from this awareness. The transition to a more sustainable kind of agriculture, in fact, can only take place through the gradual adoption of crop models that are increasingly able to produce healthy food of good quality and have access to the global commercial channels. These crop models must be in balance with the natural environment (thanks to appropriate profiles of productive efficiency), able to withstand the impact of climate change effects and harmonious with respect to the social contexts in which they should contribute to sustainable development. Within this paper, we have described the different production models and the various options available, in light of their social and economic implications. In an effort to not limit ourselves to a simple description of reality, we used a model to simulate the impact of changes in agricultural practices on the current amount of food available worldwide. It was thus possible to envisage different scenarios, assigning different values to the variables in play. The simulations carried out have confirmed the fragility of the global agricultural system and the urgency of corrective action. In particular, the search for approaches based on solutions in order to reduce energy consumption and with an in-

References of research conducted by the Barilla Group will also be noted within this paper. There was no intention to use it as an opportunity for promoting the company – the Barilla Center for Food & Nutrition exists completely independently of the Barilla Group –, but rather, as a demonstration of what a company can actually do to help solve problems. For us, this experience was the confirmation of how significant, for the whole system of stakeholders, the impact of activities to generate sustainable and widespread economic value can be.

depth knowledge of the content, will be prerequisites for sustainability.

Enjoy the read, Guido Barilla

/wido Bazica





# THE BARILLA CENTER FOR FOOD & NUTRITION

he Barilla Center for Food & Nutrition (BCFN) is a center of multidisciplinary analysis and proposals which aims to explore the major issues related to food and nutrition on a global scale.

Created in 2009, BCFN intends to listen to the demands emerging from society today by gathering experience and qualified expertise on a worldwide level and promoting a continuous and open dialogue. The complexity of the phenomena under investigation has made it necessary to adopt a methodology that goes beyond the boundaries of different disciplines.

These topics under study are broken down into four areas: Sustainable Growth for Food, Food for Health, Food for All and Food for Culture. The areas of analysis involve science, the environment, culture and the economy; within these areas, BCFN explores topics of interest, suggesting proposals to meet the food challenges of the future.

#### FOOD FOR SUSTAINABLE GROWTH

In the field of *Food for Sustainable Growth*, the Barilla Center for Food & Nutrition focuses on the issue of the optimization of natural resources within the framework of the food and agricultural sector. More specifically, the studies conducted so far have identified some critical issues and have evaluated the environmental impact of food production and consumption, putting forward a series of proposals and recommendations for individual and collective lifestyles which may have a positive effect on the environment and on natural resources.



#### FOOD FOR HEALTH

In the field of Food for Health, Barilla Center for Food & Nutrition

has decided to start its research work by analyzing the existing relationship between nutrition and health. It has studied in depth the recommendations provided by the most distinguished nutrition institutes in the world and the results of ad hoc panel discussions with some of the most accredited scientists at the international level. As a result, it has been able to provide civil society with a clear set of concrete proposals for more easily adopting a correct lifestyle and a healthy diet.





#### **FOOD FOR ALL**

In the field of *Food for All*, the Barilla Center for Food & Nutrition deals with the issue of food accessibility and malnutrition with the aim to reflect how to promote better governance of the food and agricultural sector on a global scale, in order to have a more equitable distribution of food and a better impact on social well-being, health and the environment.



#### FOOD FOR CULTURE

In the *Food for Culture* area, the Barilla Center for Food & Nutrition aims the relationship between man and food. In particular, BCFN has traced the most significant stages in the evolution of the man-food relationship, refocusing on the fundamental role of the Mediterranean diet.

In line with this approach, the activities of BCFN are guided by the Advisory Board, a body composed of experts from different but complementary sectors, which makes proposals, analyzes and develops the themes, and then drafts concrete recommendations.

One or more advisors have been individuated for each specific area: Barbara Buchner (expert on energy, climate change and the environment) and John Reilly (economist and expert on environmental issues) for the area *Food for Sustainable Growth*; Mario Monti (economist) for the area *Food For All*; Umberto Veronesi (oncologist), Gabriele Riccardi (nutritionist) and Camillo Ricordi (immunologist) for the area *Food for Health* and Claude Fischler (sociologist) for the area *Food for Culture*.

New models for sustainable agriculture constitutes the third step in a process initiated by the Barilla Center for Food & Nutrition with the papers *Is GMO agriculture sustainable?* (2010) and *Beyond GMOS: The agri-food biotechnologies* (2011). We began with the analysis of agri-food biotechnology, which has long been the focus of much debate, and then broadened the scope of the survey to the main characteristics of the different existing agricultural models in order to evaluate the profiles (current and prospective) of sustainability.

This paper attempts to investigate a key issue, especially for the future: the identification of agricultural practices and models that are truly sustainable, according to a holistic and multifunctional interpretation of "sustainability" in agriculture.

Sustainability in the food sector is, and will be, central in the near future, not only in developed countries – which, after years of intensive agriculture, are now facing the possible risks associated with energy crises and the scarcity of soil resources – but also in developing countries by which adopting less invasive modern agricultural models, could see an increase in the output and quality of their crops in a long-term vision.

Therefore, the objective of the document is to identify and examine the main factors underlying these dynamics by analyzing the most recent contributions of the scientific and institutional world, the case studies, and most interesting *best practices* worldwide in an effort to contribute to the current debate – which is fully implemented in the European Community – and offer analysis, reflection and macro policy directions in this regard.





# INDEX

Executive Summary	14
L. The future of agriculture and sustainability  BOX Food, agriculture and the scarcity of natural resources	19 22
Some data on malnutrition and undernourishment	25
2. Agriculture today: the main "agricultural models"	27
2.1 An outline of interpretation	28
Box Agricultural models and good agricultural practices 2.2 The main agricultural models identified today:	30
from taxonomy to their practical application	31
The case of Oaxaca (Mexico)	32
BOX Zero-tillage in northen Kazakhstan	35
The case of Democratic People's Republic of Korea (DPRK)	36
BOX The case of New Zeeland	39
The case of Earthbound Farms in Carmel (California, USA)	41
The case of Tigray (Ethiopia)	43
3. Sustainability of cropping systems with durum wheat in Italy:	
he case of Barilla	45
BOX The selected indicators	49
Box Barilla's Ten Commandment for the sustainable cultivation	
of quality durum wheat	55
The result of a study on durum wheat in Canada	56
4. A model of analysis and simulation	59
IDM The future of agriculture	64
5. Concluding reflections	69
Appendix. Hypothesis, assumptions and intermediate results	
of the BCFN-Millennium Institute simulation model	75
Notes and references	93



# Iture 1

## **EXECUTIVE SUMMARY**

he complexity of the agricultural system requires considering a significant number of variables that, directly and indirectly, affect the results of agricultural in terms efficiency and sustainability.

Alongside the system of food production there are fundamental aspects concerning energy, soil quality, availability/use of water resources, (agro-)biodiversity and socio-economic effects that impact agriculture at the local level. The collective impact of migration, population and different agricultural models on food security and human health are particularly relevant. Dietary habits and the consequences of climate change must also be taken into account among the major "underlying" issues in the assessment of agricultural systems.

Agriculture, in all its complexity, demonstrates daily its fragility and its exposure to possible shocks that might occur at the expense of one or more of its constituent factors; therefore, it must find new forms of balance that would allow it to be sustainable in the long run.

Achieving a more sustainable agriculture will occur through the gradual adoption of agricultural models that must be: able to produce healthy food of good quality; able to access the global commercial channels; "in-balance" with respect to the natural environment (thanks to appropriate profiles of productive efficiency); able to withstand the impact of climate change; and, finally, intune with the harmonic social contexts in which they should contribute to sustainable development.

In general, alternative approaches to agriculture can be represented in different ways, but more or less there are three fundamental aspects: economic and commercial, technological, and environmental.

According to the FAO, the systems of agricultural production can be divided into three main categories: High External Input (HEI) systems, Intermediate External Input (IEI) systems and Low External Input (LEI) systems. What is important, to this end, is the intensity of non-renewable resources consumed.

In an effort to not just give a simple description of the existing reality, but to interpret the types of current agricultural models and try to propose alternatives for the future, the Barilla Center for Food & Nutrition has made – in collaboration with the Millennium Institute – a model simulating the impact of changes in agricultural practices on the current amount of food available worldwide.

What impact could large external shocks have on world agriculture and its evolution? And, in particular, what are the effects in terms of the number of calories per capita per year available worldwide?

The proposed model allowed the simulation of different scenarios, assuming an availability of abundant energy or hypothesizing a future rapid increase in oil prices, which

would be followed by a consequent significant growth in prices of inorganic fertilizers (and therefore, a reduction in their use).

The following three macro scenarios of the global agricultural system's evolution were tested to understand implications about its future:

- → Scenario of Business As Usual (BAU): agricultural practices with a high level of external input will cover 60% of the global area/land cultivated in 2050;
- → Scenario of Strong HEI Growth: agricultural practices with a high level of external input spread at an accelerated pace to cover 90% of the total cultivated area in 2050;
- → Scenario of Stopped HEI Growth: there will be little spreading of the high external input, which will remain at the current level of 45% of the total cultivated land in 2050.

#### Results

Assuming a constant energy supply over the 80-year period observed, the production scenario with the highest yield – in terms of sustainability – was the Strong HEI Growth scenario, followed by the Business As Usual scenario (BAU) and, finally, by the Stopped HEI Growth model. In a context of simplified global development, in which possible reductions in the availability of all the elements that make up the profile of sustainability are not taken into consideration, nor any possible energy shocks, a pro Strong HEI Growth policy would generate a total availability of calories well above the required amount.

However, the Stopped HEI Growth scenario is also projected to be able to supply a more than adequate total caloric intake. This indicates that it seems there will be no problem of availability of total calories in the future.

Nevertheless, the assumption of a constant availability of energy over time is unrealistic in any case: fossil fuels are steadily decreasing and renewable energy sources are not yet a viable alternative. Therefore, it is plausible to assume that there might be a shock in the global energy supply at some point, which would put a strain on energy-intensive systems such as the HEI models.

These models become economically unsustainable and unprofitable, and serious problems would arise linked to the shift to more energy efficient models. The costs of the production change would be felt in terms of reduced output available and time spent in the acquisition of the know-how necessary for the transition.

The simulation results show that, in the case of reductions in energy availability starting in 2025, an approach to low external input would lead to a result of Worse-Before-Better (WBB), i.e., low productivity in the short-term with a return to higher levels of yield in the medium to long-term.

In the case of an energy crisis, the results depend strongly on the amount of time spent in shifting from HEI systems to a LEI (in the direction of a Stopped HEI Growth scenario) one. The results of Strong HEI Growth-Energy Shock and BAU-Energy Shock systems are less negative in the short-term.

Our simulation shows the fragility of the global agricultural system. A fragility we must deal with by promoting a balanced mix of agricultural models, built to cope with the phenomena of relative scarcity.

The reality, of course is much more complex than has been willfully represented. In addition to possible energy shocks, in fact, there are many other long-term risk factors: water availability, adaptation to atmospheric phenomena, etc.

However, the result strongly emphasizes one of the most relevant topics concerning the future: the search for knowledge-based, energy-reducing solutions will become one of the most crucial aspects of sustainability.

In the paper, there are some concrete examples of the application of the different agricultural models, also including the results of a study conducted by the Barilla Group to identify how to improve the sustainability of the main cereal crop used, that of durum wheat.

#### Conclusions

In light of the analyses, simulations and discussions carried out, what - overall – are the conclusions? The complexity of agriculture does not allow us to draw unequivocal conclusions. Nevertheless, evidence, reflections and trends emerge and converge towards a possible practical approach to true sustainability.

In particular, there are seven points of attention that we consider fundamental:

1 Sustainable agriculture is characterized by a systemic conceptual and operational approach.

For a future of sustainability, we must increasingly learn to "hold together," according to a multidisciplinary approach, the social, environmental, economic, and research and development aspects. Approaches aiming at pursuing partial goals, albeit in a very effective way, can at most reap some short-term success of one of the dimensions, but they do not help in winning the challenge of sustainability.

2 Sustainable agriculture is based on a large number of agricultural practices that are already known.

The knowledge available, comprised of scientific knowledge and proven practices, has crystallized into some excellent and practical guiding principles of truly sustainable farming. In short, these guidelines are to: cultivate a wider range of plant species, starting with a systematic use of crop rotations; minimize the mechanical working of the land; and maintain a protective cover on the surface of the organic soil. These practices/ techniques — associated with the use of high-performance crop varieties, optimized use of organic and inorganic fertilizers, integrated management of pests and diseases through appropriate practices and, when necessary, the efficient management of water resources — allow, for the same macro reference model (HEI, LEI, IEI), the achievement of better performance in terms of sustainability.

**3** Agronomic "knowledge" is not very widespread.

In the agricultural field, over the years, a wealth of available knowledge of extraordinary value has been accumulated that is only partially used today. In certain circumstances, this seems to happen due to a lack of effective processes of the transfer of know-how; in others, because it is believed that, at least partially, the technology available makes in-depth knowledge of the natural dynamics superfluous. What emerges is the need to strengthen the base of human capital in agriculture, bridging the gap between available knowledge and individual skills and of the system. Regarding this aspect, we will need to create significant investment plans because it is the prerequisite for any development in the direction of greater sustainability.

4 Correct agricultural models for specific contexts: the objective is to reduce external inputs. In our opinion, there are no good or bad agricultural paradigms a priori. There are certainly HEI models which we believe will prove to be unsustainable in fact, and LEI models which cannot be implemented in all contexts. Alongside these, there is a wide range of realities, namely that of Intermediate External Input (IEI) systems adjacent to LEI systems that can be properly managed, in light of the above sustainability requirements. In other words, what matters is the trend line, i.e., the shift toward the more sustainable IEI models and the balance between models within macro regions.

Developing countries, instead, need to adapt and revisit models that are appropriate to the specific characteristics of the local reality.

**5** Biodiversity as a tool for proper risk management.

A pragmatic approach, without prejudice, to the choice between agricultural models allows – at the level of policymaking – the maximization of the overall resilience of agricultural systems. Proper management of biodiversity and the coexistence of different models, all equally optimized as to sustainability, amplify the possibilities for responding to adverse events and specific objectives of the research system when there are alternatives (such as high quality vs. high volume).

6 Investments in technology to make agriculture more able to adapt to change.

Technology, in our view, also takes on a different connotation from the one that too often prevails. Today, in fact, when it comes to technology in agriculture, often it is referred only to the issue of productivity and yields, in the belief that they can be increased by improving the individual varieties. However, the fundamental ability is to adapt, which is expressed in the integrated and harmonious management of a wide range of tools and management logic: plant varieties that are resistant to stress, management of advanced systems of irrigation, the scientific approach to fertilization, etc.

**7** The external factors of sustainability in agriculture: food waste and losses, biofuels.

We must not forget that a large part of the problems facing the agricultural system and food are extraneous to the choices of agricultural models and their research and optimization. Some phenomena combine to emphasize the centrality of production volumes at the expense of an altogether more balanced approach: this is, above all, a matter of food waste, which has reached truly disturbing proportions and represents a challenge for future agricultural sustainability. Along with this, there emerges a central issue for the purposes of resource allocation decisions in the agricultural sector (both financial and physical): the production of biofuels. As much for the "waste" issue as for that of "biofuel," inadequate management of the problem, on the one hand, and questionable choices in the field of energy policy, on the other, result in strong pressure placed on the system of agriculture to make up for deficiencies that it should not have to bear.



# 1. THE FUTURE OF AGRICULTURE AND SUSTAINABILITY

SUSTAINABLE AGRICULTURE CAN BE DEFINED AS THE PRODUCTION OF FOOD THAT MAKES THE BEST USE OF THE GOODS AND SERVICES OF NATURE WITHOUT CAUSING

he Future of Agriculture and Sustainability is the third part of a series of position papers initiated by the Barilla Center for Food & Nutrition.<sup>1</sup> The first papers evaluated Is GMO agriculture sustainable? (2010) and Beyond GMOs. The agrifood biotechnologies (2011) respectively. Within this paper, we not only analyze agri-food biotechnology, but also different production models and their respective attributes in order to assess their sustainability.

The analysis of the agricultural system requires taking into consideration a significant number of variables that, directly or indirectly, affect the results in terms of agricultural efficiency and sustainability.


Alongside the system of food production, there are fundamental aspects concerning energy (use/production of energy, particularly from fossil fuels), soil quality (loss/depletion), availability/use of water resources (water scarcity and its use), (agri-)biodiversity and socioeconomic effects that impact agriculture at the local level. The collective impact of migration (especially in the most critical socio-economic backgrounds), population and different agricultural models on food security and human health (epidemics, undernourishment, malnutrition) are particularly relevant. Finally, the assessment of agricultural systems should take into consideration eating habits (current and future, Western and non-Western) and the consequences of climate change (the rise in average temperatures, changes in precipitation, extreme events, etc.).

The collective interaction of these variables describe an articulated and complex phenomenon - that of agriculture - which demonstrates its fragility daily. An example of this is the unresolved problem of access to food, which is expected to worsen due to the reduction of arable land, pollution and the erosion of genetic resources - and which, because of possible shocks that might occur at the expense of one or more of its constituent factors, will have to find new forms of equilibrium to be sustainable in the long run.

In light of such complexity, sustainable agriculture can be defined as "the production of food that makes the best use of the goods and services of nature, without damaging it." So, as the FAO reminds us, it should "help to conserve natural resources, to assist in protecting the environment, be appropriate for the frame of reference – from the point of view of the techniques used – and, finally, be economically and socially acceptable."<sup>3</sup>

The reasons for the growing interest in more sustainable forms of agriculture, as compared to the models that are prevalent today, lie in the increased awareness of the environmental impact of farming, as well as the awareness concerning the possible scarcity (not only in the future) of the resources that have supported the development of agriculture so far, starting with oil.

Figure 1.1. The model developed by the IAASTD to represent the complex system of agriculture (S = Same; O = Opposite; R = Reinforcing; B = Balancing)




Source: IAASTD, 2011 (presented to the Advisory Board of BCFN on February 17, 2011).

The last 50 years have been characterized by the rapid development of agricultural activity - albeit asymmetrically between the different areas of the world - toward the adoption of technologies that increase the productivity of the factors employed and a general modernization of production techniques. In some geographical areas, ever since the Sixties and Seventies, the simultaneous introduction of plant varieties that are highly responsive to production inputs (High-yielding Varieties, HYV,), the practice of monoculture, widespread mechanization, agrochemicals (massive use of pesticides, herbicides, fungicides and synthetic fertilizers developed through the use of nitrogen, phosphorus and potassium) have contributed to an extraordinary increase; at least in the short to medium-term, in production volumes for the same person – especially with regard to wheat, corn and rice, as well as generating many economies of scale along the whole chain.

On one hand, this model has ushered in a long period of high productivity and low food prices, but on the other hand, it has resulted in an intensive and often irreversible exploitation of the natural resources as the IAASTD's report Agriculture at a Crossroads reminds us: soil erosion, water contamination, pollution of rivers and water basins, deforestation and the loss of biodiversity. For this reason, over the past decade the growth of agricultural productivity has significantly dwindled, to a phase of "stagnation of yields". This is evident if one analyzes trends in the yield per hectare of corn and wheat in the U.S. (Figure 1.2.).

In the meantime, criticism of the model of intensive monoculture has led to experimentation with approaches that are more attentive to overall sustainability.




### Food, agriculture and the scarcity of natural resources

exploitation of land, particularly since the mid-twentieth century, has meant that the agricultural and food sector has been - the intense exploitation of fishing areas: ena, such as:

- the serious deterioration of arable land: 40% of the land is degraded or poor:
- tropical and subtropical forests and 45% of temperate forests have been con- - the strong dependence on fossil fuels as verted to land used for crops, including tares of peat forests in Southeast Asia,
- greenhouse gases:
- ply: by now about 70% is used;
  - an 80% use of all phosphorus available, three main producing countries;
  - an input (e.g., for the production of fer-



Figure 1.2. Trend of the yield per hectare of corn - USA (tons per hectare, 1961-2009)



Note: the yield per hectare is calculated as the ratio between the level of production and the area harvested for each year considered; the trend has been identified by using a 5-year moving average.

Source: reprocessing of data from the United States Department of Agriculture Database, 2010.

That is why the debate on the process of rethinking the logic of the prevailing models is so heated. With this in mind, it seems essential, first of all, to identify what the basic requirements are of the possible agricultural models and the issues to consider, in light of the sustainability requirements, namely:


- → The issue of controlling crop diseases and pests for ensuring adequate yields, the stability of production and food security - will still be central in the future and will also, to a certain extent - with regard to diffusion and intensity - represent a side effect of industrial agriculture.
- → What strongly emerges is the need to identify techniques and approaches to address the changes taking place (and expected to increase) in relation to two key factors: the availability of water and the quality of the soil.
- → It will also still be important to address the issue of agricultural productivity: if it is true that the problems of access to food are more related to the distribution of the product than to any insufficiency in the volume of global agricultural production,<sup>5</sup> then it is equally clear that in some parts of the world, agricultural yields still constitute a serious problem, reaching levels below those already experienced, even in the most economically advanced countries. In this sense, the correct application of agricultural techniques aimed at improving yields, remains at the heart of the debate on innovation in agriculture, especially if looking at the area of the world that requires significant improvement in average living conditions. One of the areas of greatest interest is that of finding an effective combination of the use of advanced breeding tools (MAS, TILLING, etc.) and the implementation of assessment processes and improved farming techniques and crop management.<sup>6</sup>
- → Directly and indirectly linked to all the issues listed above is a main problem related to global food security: the nutritional quality of food products. The current situation presents one of the most striking imbalances. While the number of people who are obese is increasing (especially in developed countries), there is the unsolved problem of entire populations that are malnourished (especially in developing countries), with serious implications in terms of the lack of macro- and micro-nutrients essential for a

THE GROWTH TREND OF AGRICULTURAL **PRODUCTIVITY WAS** SIGNIFICANTLY REDUCED UP TO A PHASE OF "STAGNATION OF YIELDS"

- healthy life, or even for survival. Agriculture also plays a decisive role in these issues.
- → Another fundamental issue is that of resilience: in the coming decades, it is expected that climate changes will cause structural changes in the ecological conditions of entire macro-regions of the planet, as well as a growing number of environmental shocks. At that point, it will become necessary to use agricultural and risk management techniques that are able to successfully deal with emergencies.
- → Equally crucial is the problem of food waste and food losses which if mitigated would help to fight hunger, improve food security in poorer countries and increase the income of farmers and of the consumers themselves, with positive impacts on the environment, thus avoiding loss of land, water and energy.
- → Along with the issue of the loss/waste produced by world agriculture today, what emerges - with a significant impact in terms of future agricultural sustainability - is the issue of the use of resources (financial and physical) in the agricultural sector for the production of biofuels.
- → Finally, the socio-economic aspects of sustainability must also be taken into account. Agriculture is an essential economic activity in many regions of the world, and the structure of production processes has a significant impact on the economic and social reality in which they take place. In particular, "economic and social sustainability" means, first of all, safeguarding and creating jobs in local agriculture and the improvement of living conditions in rural areas. Rural development, especially of small farmers in developing countries, should be a priority at the global level.

To summarize what has been expressed, we can say that when we use the concept of "sustainable agricultural models," we are referring to models of farming for producing healthy food of good quality, with requirements that will allow access to global marketing channels, "in-balance" with respect to the natural environment (thanks to appropriate profiles of productive efficiency), able to withstand the impact of the effects of climate change and in harmony with the social contexts in which they must contribute to sustainable development.

Figure 1.3. Food waste & losses (kg/per capita/year) in the stages of consumption and preconsumption, for the various areas



Source: Global Food Losses and Food Waste, FAO, 2011.

### Some data on malnutrition and undernourishment

world today who suffer from hunger. Out are, in fact, directly related to diseases ple, the problem of malnutrition in the laria that, in the pres ence of anapprophenomenon, 98% of the total, are contimated that 684,000 child deaths could





## 2.1 AN OUTLINE OF INTERPRETATION

IT IS POSSIBLE TO REPRESENT THE VARIOUS ALTERNATIVE APPROACHES TO AGRICULTURE BY BASICALLY REFERRING TO THREE ORDERS OF ISSUES: ECONOMIC AND COMMERCIAL ASPECTS, TECHNOLOGY AND **SUSTAINABILITY** 


he most common agricultural model of our time has been defined with different termsconventional, modern, high-input and industrial – and is considered to be the evolution of agriculture in that it includes sophisticated technologies that have greatly increased the productivity of labor.

In the Sixties and Seventies, this model led to dramatic increases in production - particularly of corn, rice and wheat - through the introduction of high-yielding varieties (HYV), monocultures, widespread mechanization and the use of agrochemicals (pesticides, herbicides and fungicides). The period in which it was adopted and developed became known as the "Green Revolution" (GR) and subsequently it was adopted in emerging contexts, such as Latin America and Asia. The practice of monoculture allows the farmer to specialize as to the factors of production used, adopting specific machines and agrochemicals and using them on many acres of land at once, thus increasing efficiency. Potentially, this leads to an economy of scale. Furthermore, the development of synthetic fertilizers formulated to provide the crops with nitrogen, phosphorus and potassium for optimal growth has helped to increase crop productivity. Therefore, the modern industrial agriculture and GR practices have resulted in the doubling of cereal production worldwide and the beginning of the surplus era.

In general, it is possible to represent the various alternative approaches to agriculture in different ways, even if all the descriptions - more or less concise - are basically made up of three orders of aspects: economic-commercial, technological and those concerning sustainability. Among the different classifications proposed in the literature, of considerable interest is the one proposed by the FAO, according to which the many farming systems can be divided into three main categories:1 the High External Input systems (HEI), the Intermediate External Input systems (IEI) and the Low External Input systems (LEI). What is particularly significant in this setting is the reference to the intensity of the non-renewable resources consumed. Figure 2.1. shows the conformation of different production systems, depending on the degree of substitution between processes primarily based on inputs consisting of natural resources and processes mainly based on synthetic or technological inputs.

The HEI systems (High External Input) are characterized by their strong commercial orientation, the use of crop varieties with high yield, intense mechanization (which is accompanied by low labor intensity) and dependence on productive factors of a synthetic nature (fertilizers and agrochemicals). These models are aimed at maximizing production output in conditions of maximum efficiency due to achievable economies of scale. Monoculture and the cultivation of crops of genetically modified organisms (GMOs) represent the extremes of this approach. At the other end of the spectrum lie the LEI systems (Low External Input) which are characterized by the use of traditional plant varieties, the use of labor-intensive techniques and knowledge, and the low use of chemicals.

Figure 2.1. The three main agricultural models according to the FAO



Source: FAO/OECD, Food availability and natural resource use in a green economy context, 2011.

Most agricultural models can be found at an intermediate level, with a range of possi-

These are the so-called IEI models (Intermediate External Input), involving the use of improved crop varieties through conventional breeding techniques and hybridization, the search for a sustainable balance between mechanization and labor, the use of techniques with a high content of knowledge and the use of fertilizers and chemicals, according to integrated patterns of cultivation.

The sustainability of the different models is obviously different: in particular, the HEI systems seem to be able to ensure crop yields that are better in terms of product per area, but they are obtained with a high consumption of resources, which undoubtedly makes them more fragile as to the future prospect of a possible scarcity of resources. The LEI systems, however, are forced to "pay" for their minimal impact on resources in terms of crop yields that are typically lower. In any case, this is a very simplified representation of the reality, which instead, is far more complex and dynamic. Therefore, it seems useful to briefly analyze the characteristics and specific cases of the application of some of the main agricultural models worldwide, taking into consideration the effects in terms of sustainability.

In particular, sustainable agricultural practices include: the promotion of biodiversity, the recycling of plant nutrients (nutrient cycling), protection against soil erosion, the conservation and protection of water, the minimum processing of the terrain, the absence of chemical products and synthetic fertilizers, and integration between agriculture and raising animals.

## Agricultural models and good agricultural practices

The description of the possible models of agriculture according to a tripartite classificathat actually exist in the world. This simplified representation has been used to facilitate the consideration of some important trends. In

- models" should start from the definition of typical situations regarding the use of key inputs, such as:
- modern varieties obtained by conventional breeding, hybrids and GMOs;
- water resources: rain-fed cultivation and crop irrigation;
- energy resources: labor intensive and highly
- soil resources (quantity): extensive farming
- soil resources (quality): zero/minimum tillage and conventional plowing:
- biological control, integrated pest management and the fight against synthetic che-
- → Each form of agriculture (or agricultural stract, but must always be placed in a geographical context with regard to the climate, soil, and economic and social development.
- with the development of growing uncertainties about the factors that define the

models can provide an essential model of risk management. A main model can a smaller scale, to be rapidly expandable high inputs, while, however, at the same ties of traditional corn to be used when of fertilizer and energy, etc.).

- ally plan the global agricultural producresources and access to food are ensured able land in order to optimize the use of the scarce basic resources. Indeed, it is to produce rain-fed cereals in temperate regions of the planet than to pursue the self-sufficiency of individual countries.
- model) can never be assessed in the ab- > The matter of "good agricultural pracways, they can be applied to each type same is true for all the other "best prac-

# 2.2 THE MAIN AGRICULTURAL MODELS IDENTIFIED TODAY: FROM TAXONOMY TO THEIR PRACTICAL APPLICATION

#### Traditional Agriculture

Traditional agriculture includes forms of breeding arising from the co-evolution of local social and environmental systems. It has a high ecological logic expressed through the intensive use of local knowledge and natural resources, including the management of agrobiodiversity as a form of a diversified agricultural system.



James P. Blair/National Geographic Stock

#### The case of Oaxaca (Mexico)

The CEDICAM (Center for Integral De- nation also occurred in the criollo corn velopment of the Mixteca Alta for farm- varieties from the GMO varieties, which and a group of farmers in Mixteca for practices that had been carried out for the purpose of creating a democratic thousands of years. organization that would promote sus- It was then that the CEDICAM began its community projects. The Center was es- the local ecosystem through the adoptablished to find a solution to the wide- tion of sustainable agriculture. spread environmental degrad ation in In 2008, Leon was awarded the Goldman the region, which had been taking place Environmental Prize for his pioneering about five hundred years ago.

to North America in search of work.

to NAFTA), it became clear that this new However, it should be emphasized that

out under the CEDICAM. He managed to In fact, because of the large-scale de- unite farmers who, together, planted one forestation carried out to allow for the million native trees in the Mixteca region one of the most eroded in the world. Ac- cal biodiversity. Leon also turned to the amounting to 500,000 acres of land. To canals were built by a small number of complicate the situation, due to NAFTA peasants, who thus also increased their longer able to earn a living only with the canals that kept the water in situ, allowvived and animals have begun to reap-At first, it was thought that a solution pear. Leon has also promoted the use of could be the implementation of modern greenhouses to produce vegetables for by the inhabitants of the region who concerning sustainable/basic agricultural

system of agriculture, in fact, was not an important factor for the success of would only further erode the soil and put once again was certainly the fact that the the farmers into debt. Moreover, in the peasants of CEDICAM understood that outside are not always the best for the er" in order to allow people to address to work again with their families.

than foreign products. And as a result of arose from a local need, was developed the work of CEDICAM, thanks to the eco- by the local people and focused on real nomic improvement of the area, some of issues affecting the region. The locals the people of Oaxaca who had emigrated know their own land and traditions betto the United States have returned today ter than anyone else and are, therefore, more effective partners to promote the The CEDICAM is an example of a move- development of social capital and food



#### Conservation Agriculture

CONSERVATION AGRICULTURE IS A MODEL THAT PROMOTES LITTLE OR NO WORKING OF THE SOIL SO AS TO MAINTAIN AND PRESERVE ITS STRUCTURE

Conservation agriculture (CA) is a model that promotes minimal or no working of the soil in order to maintain and preserve its structure; in addition, it uses practices such as mulching and does not limit the use of GMO seeds or agrochemical products, as is the case with organic farming. The FAO promotes conservation agriculture in developing countries and defines it as an agricultural model that "aims to support sustainable and profitable agriculture and, consequently, its objective is the improvement of the living conditions of the farmers through the implementation of three principles: minimum tillage, mulching and crop rotation".

Both conservation agriculture and that of zero-tillage involve a minimal processing of the soil, planting in rows and the maintainance of crop residues. The proponents of conservation agriculture wish to preserve the soil's micro-organisms living in the surface soil and assist the plant in the absorption of water and nutrients. The zero-tillage also allows for maintaining the moisture of the soil, thus reducing water usage and improving the absorption and retention of CO<sub>2</sub>.

According to Rasha Omar of the IFAD, about 95 million hectares around the world are managed through conservation agriculture: of these 25% is located in the United States, approximately 23% in Brazil and 18% in Argentina (Derpsch, 2005 - quoted by the IFAD) and in all three cases, these are almost always crops (mainly soybeans) that come from herbicide-tolerant GMOs.



© Corbis

## Zero-tillage in northern Kazakhstan

No-tillage (or zero-tillage) is a technique horizon of the root system and thus preused in a highly conservative concept of venting soil erosion. The snow is retained sion in northern Kazakhstan, so much so left on the ground, even up to a height that from 2007-2008, it was estimated of about 35-40 cm. (Fileccia, 2009). The that the area under zero-tillage had dou- lack of cultivation of the soil causes a rebled, reaching up to 1.2 million hectares. duction in costs, but this savings is often has been a decline of traditional plowing, purchase of herbicides, since it has been while techniques of little or no tillage have shown that the zero-tillage farming is are beginning to see a 20-50% increase pletely eliminated (Fileccia, 2009).

The zero-tillage farmers use special ma- the high cost of specialized machinery, ground. These two methods have proven for their towing), but in fact, many farms very effective in maintaining soil mois- converted to zero-tillage in Kazakhstan for 35-40% of the annual rainfall and is national community for its development, an important part of the water supply in Kazakhstan has become one of the top Its slow and steady melting allows the extension of areas cultivated with the

## The case of Democratic People's Republic of Korea (DPRK)

From September 2002 until November leguminous crops in order to select those 2005, the FAO carried out a project on which were the most suitable to the real-

curity aimed at obtaining food security The introduction of conservation agriculfor the Democratic People's Republic of ture practices in North Korea, for example, Korea. About 80% of the population of brought about a reduction in the number livelihood is closely linked to agriculture thanks to the mulching. There were also (FAO, 2007), but due to the harsh climate increases in yield compared to traditional with temperatures ranging from -19°C work systems of 0.41 to 0.63 t/ha (FAO, in winter to 25°C in summer, the period 2007). However, in 2003, areas under years (FAO, 2007). And finally, the soils companies under study reported no inended up typically poor in organic matter crease in production, because it had not aged the adoption of cropping systems quently, the cover crop grew in such guanthat were an alternative to the traditional tities as to compete with the corn, thereby

caused by traditional plowing (20 cm). increase of nutrients and organic mat-The objective of the FAO project in North ter as compared to that of conservation ple, should have been replaced by no-till mg/kg of soil and the available phosphotechnologies and cropping systems that, rus by 30-40 mg/kg of soil (FAO, 2007). among other things, included green ma- In land cultivated with a conservation was conducted on plots of 50 acres be- ture was also found and it has been estilonging to three cooperatives; economic mated that the mulching with straw was facilities were used for direct seeding de- able to increase the moisture content by tion of wheat-soybean, corn-soybean and protection from the erosive action of water.

formed to determine if conservation ag- age saving of 15.5 kg of fuel per hectare.

consumption and labor hours required be a valid alternative, both economically per hectare, it was estimated that con- and ecologically sustainable, to convenservation agriculture practices allowed for tional agriculture. As a result, farmers are duction factors (FAO, 2007). In addition, agriculture in their region and fifteen forusing conservation agriculture practices, eign companies have expressed interest



#### Biodynamic Agriculture

The founder of biodynamic agriculture, Rudolf Steiner, spoke of the biodynamic agricultural model for the first time in a series of conferences held in 1924. Since then, the practice has spread throughout the world to include over 4200 biodynamic farms, with 128,000 hectares of land, in 43 countries (Turimek, 2009).

Biodynamic farming has the non-use of chemicals and synthetic fertilizers in common with organic farming, but instead uses manure as fertilizer, then calls for the rotation of crops, pest control carried out in a natural way, and diversification of crops and livestock.

However, biodynamic farming differs from organic farming and other agricultural models in the preparation of compost and plant protection products. There are eight preparations: 500, 501, 502-507 and 508, corresponding to cattle manure, silica, yarrow flowers, chamomile, dandelion, valerian, oak bark and nettle. Preparations 502-507 are added to the compost, while the others are diluted in water and sprayed directly on the field.

Moreover, biodynamic farmers follow the lunar cycle for planting and sowing, convinced of the validity of the effects that the moon could have on plant growth.



© Corbis

#### The case of New Zealand

In 1993, a study was conducted in New were found to be higher in the biodynam-(topsoil) of 2.2 cm.

though the study was conducted for four agriculture. However, concentrations of

prise (crop production and livestock) and analysis showed that biodynamic farms pared to conventional agriculture farms. conventional farms (regarding vegetables alternation of nutrients in the soil. The ity and is becoming increasingly important it would have a significant impact on the

#### Industrial Biological/Organic Agriculture

AGRICULTURE AROSE IN THE UNITED STATES IN THE LATE NINETIES, WHEN THE USDA ESTABLISHED NATIONAL ORGANIC STANDARDS

THE INDUSTRIAL ORGANIC There is no official definition of industrial organic farming, but we can say that it arose in the United States in the late Nineties, when the USDA established national organic standards.

> The reason the USDA found it necessary to develop organic standards was due to the entry of large food companies within the organic market. After all, companies like Earthbound, Cascadian Farm and Horizon practice a form of agriculture that is not unlike modern industrial agriculture, capable of feeding the global food economy.

> Economies of scale are obtained with the use of mechanization and monoculture, an aspect not contemplated by the organic principles, and consequently the biodiversity is often reduced in these large organic farms. However, among the environmental benefits of this production model is a ban on use of fertilizers and pesticides, which means greater protection of land, water courses and ultimately the consumers themselves.

> Nevertheless, the sustainability of industrial organic farming has some ethically and socially negative aspects. For example, unlike organic farming, the industrial model is less tied to regionalism and practices adapted to local conditions; instead its products are shipped throughout the world (which allows Brazilian consumers to eat organic lobster from Maine), while the original organic movement promotes the purchase of local products and the strengthening of relations between communities and farmers.

> For large food businesses, less stringent national organic standards provide an opportunity to benefit from economies of scale and to take advantage of the transition to organic production. However, the entry of large companies in a growing organic market has negative implications for small farmers because they find that they have less control over the prices and are often exploited by large companies that are able to produce more with less.

> Industrial organic and biological models indeed seem to actually be two distinct models: the first is aimed at a high profit and the second has important social and ethical objectives. It would be more appropriate, therefore, to give the large organic farms, which are limited to meeting national standards, a name that is different from the one given to small organic farms, that aim to ensure food sovereignty and rural employment.

> At the beginning of the movement in the early Sixties, thousands of small organic farms were purchased by large food companies. This was the case of the Cascadian farms which, once at the forefront of the "back to earth" organic movement, are now owned by General Mills; Dagoba is now owned by Hershey's and Horizon and Alta Dena are owned by Dean. The advantage of the management buy-in of organic farming is the price reduction of many organic products, which currently represents precisely the main reason for their limited purchase by customers.

## The case of Earthbound Farms in Carmel (California, USA)

ferent areas of California, two in Arizona, to be considerable, but to accurately cal-Mexico (Shapin, 2006).

sold in the United States.

of which is 680 acres (275 hectares).

Earthbound Farms with two and a half persistent pesticides and more than Carmel, California. In 1986, they started chemical fertilizers." Moreover, they claim the retail sale of ready-made salads and to "save about 1.8 million gallons (about now have become the major producers of 7000 liters) of oil by avoiding the use of organic products and the most success- pesticides and fertilizers made from peful in the world, billing up to 450 mil- troleum, and to fight global warming by lion dollars in 2006 (Shapin, 2006). They avoiding the release of carbon dioxide, a need six companies to meet the demands major greenhouse gas, into the atmosof large retailers like Costco and Whole phere, equivalent to that produced by one in Colorado, and in three regions of culate the emissions of carbon dioxide, the gases emitted by the trucks for the Earthbound Farms has a huge influence transport of compost and the mechanized on the organic market, seeing as they pro- harvest of thirty-six thousand acres need duce more than 70% of the organic lettuce to be added to the equation. The transport chain of Earthbound products on the hectares) on its large farms, the largest son that emissions produced going from Mexico to California to New York and in

#### Organic Agriculture

IT IS BASED ON THE ECOLOGICAL PROCESSES, ON THE PRESERVATION OF BIODIVERSITY AND ON THE TRENDS OF THE PRODUCTION CYCLES UNDER LOCAL CONDITIONS

Organic agriculture has been defined by the IFOAM (International Foundation of Organic Agriculture Movements) as "a system of production that sustains the health of soils, ecosystems and people." It is based on ecological processes, on the preservation of biodiversity and adaptation to the production cycles of local conditions, rather than the use of

Organic agriculture combines tradition, innovation and science for the benefit of the environment and promotes fair relationships, as well as a good quality of life, for all involved (IAASTD, 2009).

There are many definitions of organic farming, and more are being made in every new country.<sup>2</sup> We can summarize and say that what are considered to be organic farming practices (even though adopted also in agricultural models that do not qualify as "organic") include: the use of cover crops, crop rotation, green manure, composting and the use of catch crops.

From 2007 to 2010, the amount of certified organic farmland increased by 3 million hectares, equal to 9%. In 2010, the global market for certified organic products reached \$55 billion, making it the fastest growing sector in the food business economy.

The massive increase in the popularity of organic products is mainly due to the perception of nutritional and health benefits (mostly environmental) that these bring.

The premium prices of organic products have led to the increase in the income of organic farmers, also benefiting the small landowners in developing countries.

In 2010, 37.2 million hectares of land were farmed organically, equal to an increase of 6.2% over the previous year. The countries with the largest number of organic producers are India (340,000), Uganda (180,000) and Mexico (130,000). In India, the main reason farmers choose to switch to organic production is linked to health benefits for the farmer and also for the consumer.



Anthony Stweart/National Geographic Stock

## The case of Tigray (Ethiopia)

The "Tigray Project" in Ethiopia is one of vation, the reduction of grazing and the adoption of organic agriculture can sub- woody plants, the creation of community stantially increase yields and, at the same ponds, small dams and river diversions, time, result in increased access to food by in order to collect and store water for the population groups considered to be use in the dry season, the promotion of dices of poverty (Edwards, 2007).

northern Ethiopia, where the diversity of the household and composed of elderly experts who want to find out if a com- and plants used as fodder to be sold loproduction through ecological principles education, enabling them to acquire the "Tigray Project," the Institute for Sustain- simple technologies and tools that are able Development, in collaboration with easy to find and use (such as foot pumps). tection Authority (EPA) of Ethiopia adopt- were analyzed. Overall, the use of comed the principle of Extension Services post had doubled the yields of all crops: (technique based on direct and constant beans, barley, wheat, teff, hanfets (a mixthat of the field technicians - Edwards, was also a general increase of biomass, 2010). Thanks to the success achieved although to an extent not comparable to and recognition by the government that the increase of yield. strategy against land degradation and for sons for the increase in yield is that the project has been extended to other Ethio- own seed varieties, which were therefore pian regions (Edwards, 2010). A 2010 adapted to local conditions. In addition, FAO publication of the "Tigray Project" farmers observed that the use of compost

bio-pesticides based on local knowledge; Tigray is located in the highlands of support for families with female heads of



# 3. SUSTAINABILITY OF CROPPING SYSTEMS WITH DURUM WHEAT IN ITALY: THE CASE OF BARILLA

ustainability in agriculture is one of the objectives of the Common Agricultural Policy. In this regard, sustainable agricultural systems are those production models that are able to produce foods that are adequate in quality and quantity, ensure a fair economic remuneration for the farmers and help safeguard the agricultural soil and natural resources. In other words, sustainability means "seeking a long-term maintenance of the agricultural production and soil fertility, while reducing environmental risks related to those very same agronomic practices."

But just how much impact does agricultural activity have on the entire life cycle of an agro-food product? To answer this question, the case of durum wheat semolina pasta was examined; its impacts have been analyzed through the Life Cycle Assessment (LCA), an objective method of evaluation and quantification of the environmental and energy loads and the potential impacts associated with a product/process/activity along the entire life cycle, from the acquisition of raw materials to end of life ("cradle to grave").

This analysis showed that the cultivation phase of wheat, with its agronomic practices, together with cooking, is one of the most important phases in terms of environmental impact (Figure 3.1.).

Using greenhouse gas emissions as an example, we can see that the major impacts associated with farming activities are due to the use of nitrogen fertilizers and mechanical operations, in particular for working the land.


These results, combined with a need to analyze and evaluate Italian agricultural systems where durum wheat is grown, led Barilla to launch a multidisciplinary study that simultaneously considers the economic, production, agronomic, environmental and food safety values.

The ultimate goal was to identify "sustainable" agricultural systems, validate the various areas of national production and to raise both the quality and quantity of raw material. Once validated, these systems should be introduced in the discipline of durum wheat cultivation.

From a methodological point of view, four main areas have been identified: the plains of the Lombardy-Veneto regions, Emilia-Romagna, Central Italy (Tuscany, Marche and Umbria) and Southern Italy and islands (Puglia, Basilicata and Sicily); the standard crop rotations identified are sufficiently representative of the rotations in which durum wheat is grown in Italy (Figure 3.2.).


Figure 3.1. Results of an LCA analysis of durum wheat for pasta

#### 2. Milling of wheat




#### 5. Transport

#### CO<sub>2</sub> emissions – equivalent (g CO<sub>2</sub>/500g of pasta)



\*Carbon sequestration means the amount of  $CO_2$  that was absorbed by the wheat during its growth. The value is normally shown separately from the others and not added, in that from the scientific point of view, there is no agreement on how the reporting of this data should be done.

Source: Environmental product declaration of Durum wheat semolina dried pasta produced in Italy, in a cardboard box; S-P-00217; 10/03/2011. Data regarding the average Barilla production worldwide, www.environdec.com.



Source: Sustainability of Cropping Systems with durum wheat, in "Grano Duro News," 2011.

The agronomic and economic studies were supported by the environmental assessments conducted by using the LCA methodology and synthesized through the use of the following indicators: the Water Footprint, the Ecological Footprint and the Carbon Footprint, as shown in the following table:

	SYSTEM	Grain Yield (t/ha)	Carbon Footprint (t CO <sub>2</sub> /t)	Water Footprint (m³/ha)	Ecological Footprint (gha/t)	Gross Revenue (€/t)	Nitrogen use efficiency (kg/kg)	DON Risk (0-9)
CENTRAL ITALY	Cereal*	3,3	0,67	745	0,73	24,1	28,4	3,9
	Forage	4,3	0,30	478	0,47	99,4	66,7	0,0
	Industrial	5,3	0,43	502	0,49	138,8	45,3	0,0
	Protein	5,3	0,34	479	0,47	139,2	58,5	0,0
EMILIA- ROMAGNA	Cereal*	7,3	0,51	328	0,40	140,7	32,5	7,9
	Industrial	7,5	0,41	315	0,38	156,7	42,2	2,3
	Industrial Horticulture	7,5	0,36	315	0,38	151,1	47,1	1,7
LOMBARDY VENETO REGIONS	Industrial*	7,5	0,42	294	0,36	166,9	44,0	1,7
	Corn-growing	7,0	0,51	315	0,38	155,2	33,8	7,9
SOUTHERN ITALY AND SICILY	Cereal*	2,5	0,74	1429	1,11	23,3	32,4	1,1
	Forage	5,0	0,45	694	0,54	132,8	44,3	0,0
	Industrial Horticulture	4,2	0,53	874	0,68	111,8	38,7	0,0
	Protein	5,0	0,45	694	0,54	132,8	44,3	0,0

<sup>\*</sup>Standard crop rotation normally adopted in each area.

Source: Sustainability of Cropping Systems with durum wheat, in "Grano Duro News," 2011.

#### The selected indicators

- → Wheat production: for each system analyzed, the production of different crops have been analyzed, including durum wheat. In particular, for the latter, the data was reported in tons of grain per hectare to 13% humidity. The yields reported in the paper refer to medium-high productions for every turnover and consider the use of Good Agricultural Practices (GAP) for the various production processes.
- → Carbon Footprint: represents the total amount of GHG (Greenhouse Gases), which are those substances in the atmosphere, both natural and anthropogenic, that are transparent to incoming solar radiation on Earth, but which can substantially retain the infrared radiation emitted by the surface of the Earth by the atmosphere and clouds. Greenhouse gases are expressed as mass of CO₂ equivalent, treating all gas emitted in terms of global warming effects of the CO₂, according to the conversion tables defined by the IPCC (International Panel on Climate Change). In the case of this study, the Carbon Footprint is expressed in tons of CO₂ equivalent per ton of durum wheat produced.
- Water Footprint: represents the consumption of water related to the production of goods and services. About 85% of the human Water Footprint is related to agricultural (and animal) production, 10% to industrial production and 5% to domestic consumption. In the case of this study, the indicator measures the water consumption of the cultivation of durum wheat in terms of the volume of water consumed during the various production processes and following the natural evaporation of the crops. Irrigation was not considered because it is a practice that is not usual in these distribution areas. The Water Footprint is expressed in cubic meters of water per ton of wheat produced.
- → Ecological Footprint: measures the area of biologically productive land and sea need-


- ed to regenerate the resources consumed by a human population and absorb the corresponding waste. Using the Ecological Footprint, it is possible to estimate how many "Planet Earths" would be needed to support humanity if everybody lived according to a certain lifestyle. The present study used the measurement in "global hectares" per ton of wheat produced.
- → Gross Income (GI): represents the difference between the GMP (gross marketable production, updated according to prices in March, 2011) and the Cost of Production of the crops. The GMP does not take into account the costs of the direct support and/or indirect effects of the CAP, whereas the Cost of Production takes into account only the direct costs of cultivation (cultivation operations and technical means) and not those that are indirect (land use, financial interests, taxes and fines, etc.). In the present study, Gross Income was measured in Euros per ton of wheat produced.
- → Nitrogen use efficiency (NUtE): represents the amount of grain produced per unit of nitrogen distributed on the crop of durum wheat.
- → DON Risk: expresses the risk of contamination of grain by Deoxynivalenol (DON), a dangerous mycotoxin that is developed by a group of pathogenic fungi (Fusarium spp.) that attack the ear of durum wheat. The mycotoxin risk index combines the meteorological factors conducive to the production of mycotoxins by Fusarium graminearum and F. culmorum with the predisposing factors, and the specifics of the productive unit, such as varietal susceptibility, crop precession, and the processing of soil. The mycotoxin risk index ranges from 0 (recorded when there are no conditions for the production of mycotoxins) to 9 (recorded when the conditions are very favorable for the production of mycotoxins).

#### Results of the study

Figure 3.3. (A, B, C and D) reveals graphical results of the study of the effects of cultivation systems on the Carbon Footprint, Gross Revenue, Efficiency of utilization of nitrogen and DON risk.


With regard to the Carbon Footprint (Figure 3.3. A), within each macro area we can observe an interesting variation in cereal systems: in principle the cultivation technique of durum wheat has the most impact in terms of greenhouse gas emissions. This is partly explained by the fact that in such systems, in order to grow durum wheat, cultivation re-

Figure 3.3. Effects of cropping systems on the Carbon Footprint (A), Gross revenue (B), Efficiency of the utilization of nitrogen(C) and DON Risk of durum wheat (D)




\*Standard rotation of crops normally adopted in each area.

<sup>\*\*</sup>The difference between the average values measured for crop rotations and the values recorded in the cereal systems.




\*Standard rotation of crops normally adopted in each area



\*Standard rotation of crops normally adopted in each area.

\*\*The difference between the average values measured for crop rotations and the values recorded in the cereal systems.



Source: Sustainability of Cropping Systems with durum wheat, in "Grano Duro News," 2011.

quires very expensive operations, such as plowing, in order to reduce the risk of mycotoxins or significantly increase the intake of artificial nitrogen, since cereal crops in rotation (common and durum wheat, corn and grain sorghum) remove large quantities of the element and leave crop residues that are not easily degradable by the microflora in the soil. On the other hand, especially where there are forage or protein crops in the rotation, the "environmental cost" decreases significantly. In these cases the residual nitrogen in the crop rotation makes a reduction of inputs of artificial nutrition possible, and it is possible to use conservative tillage techniques: minimum tillage or direct seeding.

The economic analysis of the cost of durum wheat also reflects the considerations mentioned above (Figure 3.3. B). The cereal systems, especially those of Central and Southern

<sup>\*\*</sup>The difference between the average values measured for crop rotations and the values recorded in the cereal systems.

Italy, are at the limit of affordability, considering that the prices of wheat applied are those of the Commodity Exchange recorded in Bologna at the time this paper was written (280 €/t).

The efficiency of nitrogen use in wheat was found to be higher in rotations in Central Italy, particularly at the end of the cycle of alfalfa, a crop that leaves large amounts of available nitrogen in the ground (Figure 3.3. C).

The DON Risk (Figure 3.3. D) was calculated by using the mathematical models from the Catholic University of Piacenza, which attribute half of the risks to climatic variables and the remaining half to agronomic factors (precession, type of tillage, variety, etc.). As is known, the risk of mycotoxin is greater in the areas of the North and in particular, in the farming systems of corn in the plains of Lombardy and Veneto, and of cereal in Emilia-Romagna. However, the risk remains, albeit at very low levels, even in the macro areas of Central and Southern Italy, where cereals are the predominant crop.



© Corbis

#### Conclusions

The study conducted showed the possibility of assessing the "sustainability" of a crop or a cropping system using multi-disciplinary analysis, combining different environmental, agronomic, economic and food security indicators. This type of holistic view of the cropping system opens important perspectives of strategy and possible scenarios in order to achieve the objectives of a modern agriculture.

It is evident that the characteristics of a species, in this case durum wheat, are strongly linked to the context (system or model) in which it is grown. Not only do all of the parameters of "sustainability" change substantially, but so do the final quality and quantity of the material produced. But the most interesting thing is to discover that the application of traditional farming practices, and a proper crop rotation in particular, ensure environmentally sustainable production.

The next step is the "actual measurement in the field" of the most favorable cropping systems, which will be compared to traditional ones.

The next step is to directly involve farmers and industry experts, so that the concepts of sustainability and productivity come to be part of corporate strategies as two absolutely reconcilable aspects of agricultural production. In order to achieve this goal, beginning with the next sowing, pilot crops will be implemented in more favorable rotation contexts which are also more economically sustainable than the ones today, in a network of farms distributed throughout Italy (Figure 3.4.).

To better explain this to farmers, Barilla developed "Ten Commandments" for the sustainable cultivation of quality durum wheat.

Figure 3.4. Project Durum wheat and Sustainable Agricultural Systems. Provision of platforms for validation of the "Pilot" companies



**ENVIRONMENTALLY** SUSTAINABLE **PRODUCTION** 

THE APPLICATION OF

PRACTICES, AND IN

PARTICULAR PROPER

CROP ROTATION, ENSURE

TRADITIONAL FARMING

Source: Sustainability of Cropping Systems with durum wheat, in "Grano Duro News," 2011.



## Barilla's Ten Commandments for the sustainable cultivation of quality durum wheat

the cultivation of durum wheat: a list of guiding principles for farmers who are tainable agriculture.

favorable crop rotations, to the efficient use 7. Restrain weeds in a timely manner: of resources and the proper use of techniboth qualitatively and quantitatively.

- cereal crops are, in fact, the cause of 9. Protect the plant from disease: carry high environmental impacts and low
- are adapted to the specific conditions, the following guidelines.
- to be sown in relation to the cultivation

- area and expectations in terms of pro-
- identity (production capacity, techno-
- thickly prevents the crop from making causes enticements.
- the treatment must be timely and apcropping practices.
- needs of the plant: the use of nitrogen
- out the treatments of defense in relacomprehensive strategy that involves all aspects of cultivation.
- rum wheat in the cropping system (rotation) without limiting it to the context




#### The results of a study on durum wheat in Canada<sup>1</sup>

Improving cropping systems may help mitigate greenhouse gas emissions. This study determined the carbon footprint of durum wheat (*Triticum turgidum L.*) produced in diverse cropping systems.

Durum was grown in rotation systems which had different combinations of oilseed, pulse (legumes) and cereal crops at five site-years in Saskachewan, Canada. Total greenhouse gas emissions from the

decomposition of crop residues along wit various production inputs were used for the estimation of the carbon footprint. Of average, the emissions from the decomposition of crop straw and roots accounted for 25% of the total emissions, those from the production, transportation, storage and delivery of fertilizers and pesticides the farm gates and their application 43%, and emissions from farming operations 32%.

Figure 3.5. Carbon Footprint of durum wheat grown under different cropping systems in southwest Saskachewan, Canada. Data was calculated from the averages of three rotation cycles conducted in fields in the period of 1996-1998 (first cycle), 1997-1999 (second cycle) and 1998-2000 (third cycle). Bars are standard errors



Source: Gan, Y., C. Liang, X. Wang, B. McConkey, *Lowering Carbon Footprint of Durum Wheat by Diversifying Cropping* Systems, in "Field Crops Research", 122 (3), pp. 199–206, Elsevier, 2011.





# 4. A MODEL OF ANALYSIS AND SIMULATION

THE AMOUNT OF FOOD PRODUCED EACH YEAR IS ENOUGH TO FEED THE WORLD, EVEN IN THE **FUTURE** 

n an attempt to interpret the types of current agricultural models and to propose alternatives for the future, the Barilla Center for Food & Nutrition has created, in collaboration with the Millennium Institute, a simulation model to study the impact of changes in current agricultural practices on the amount of food available worldwide.<sup>2</sup> The results of this model are the basis for many of the ideas proposed so far.

The objective of the analysis is to understand how far-reaching external shocks, summarized here as a very significant increase in the price of oil, can have an impact on world agriculture and its evolution, expressed in terms of the agricultural models adopted.

In particular, two main patterns of agriculture have been identified and defined:


- a Low External Input model (LEI), characterized by a low energy intake and an elevated use of labor;<sup>3</sup>
- a High External Input model (HEI), characterized by a high consumption of energy and inorganic fertilizers.4

The two models mainly differ as to the different characteristics of sustainability over time. Considering a period of 80 years (1970-2050) and evaluating the impact on the amount of calories produced per capita annually, assumptions can be made concerning the choices of the most appropriate production policy.

It is important to clarify that, starting from the simulations carried out, <sup>5</sup> the amount of food produced each year is enough to feed the world, even in the future taking into account a rate of productivity growth in line with the current demographic trends and projections made by the FAO and the OECD.6 A significant portion of the problems the food system faces – as we mentioned – depends on critical issues related to distribution, tintended use and food product waste.


- → Under conditions of abundant energy supply, the simulation model includes three different scenarios (Figure 4.1.):
- Scenario of Business As Usual (BAU): practices at a high level of external input will cover 60% of the total area cultivated in 2050;
- Scenario of Strong HEI Growth: practices with high external input will spread at an accelerated pace and will cover 90% of the total area cultivated in 2050;
- Scenario of Stopped HEI Growth: we will witness the poor dissemination of models of the high use of external inputs, which will retain their current share of 45% of the cultivated land in 2050.
- → There is also a simulation (Figure 4.2.) of a rapid increase in oil prices between 2025 and 2030 (a context defined "Very High Energy Price"): oil prices will rise rapidly in 2030 to reach \$200 a barrel, and then level off at around \$280 per barrel in 2050. Because of the rapid rise in oil prices, the price of inorganic fertilizers will grow so much as to substantially

Figure 4.1. Agricultural production for human nutrition



Source: FAO data base.

Figure 4.2. Agricultural production for human nutrition in a case of energy shock



Source: FAO data base

THE THREE SCENARIOS ANALYZED: A) BAU-ENERGY SHOCK, B) STRONG HEI GROWTH-ENERGY SHOCK AND C) STOPPED HEI GROWTH-ENERGY SHOCK reduce their use; these conditions, will benefit only the high-value crops and in general only 50% of the area cultivated globally. Therefore, the effects were estimated on the number of calories per capita per year for each of the three basic scenarios (BAU-Energy Shock, Strong HEI Growth-Energy Shock, Stopped HEI Growth-Energy Shock).

What has been made is a simulation model that tries to understand what the result of a global shock could be, depending on the configuration of the global agricultural system. The discriminant variable is the use of energy as the primary factor of production.

If we hypothesize, for example, a constant availability of energy over the observed period of 80 years, the production scenario with a higher yield – in the scheme of sustainability- would be the one of Strong HEI Growth, followed by the Business As Usual (BAU ) scenario and, finally, by the Stopped HEI Growth scenario (Figure 4.1). In a context of simplified global development, in which possible reductions in the availability of all the elements that make up the profile of sustainability are not taken into consideration, with the certainty of not being subject to energy shocks, a pro-Strong HEI Growth policy would generate a total availability of calories well above the required amount. It is interesting to note that also the Stopped HEI Growth scenario seems to be able to provide, a projected total caloric intake that is more than  $\frac{1}{2}$ adequate. This indicates that there does not seem to be an issue with the availability of total calories. However, it must be said that this simulation model does not take into account the inequalities between different geographical areas, which, in fact, constitute the real problem. Moreover, the assumption of a constant availability of energy over time is unrealistic, because fossil fuels are steadily declining and renewable energy sources are not yet a viable alternative. To conclude, it can be assumed that at some point in time, an energy shock of the global supply is likely to occur, which would put a strain on energy-intensive systems such as the HEI models. In fact, these models would become economically unsustainable and unprofitable, and serious problems related to the transition to more efficient models from the point of view of energy use would occur. The costs of the change in production would become evident in terms of the minor output available and of time spent in the acquisition of the know-how needed for the transition. Figure 4.2. shows the estimated affects of an energy shock on global energy output between 2025 and 2035.

The simulation results show how if reductions in energy availability occur starting in 2025, an approach to low external input would lead to a Worse-Before-Better (WBB) result, or low productivity in the short term with a return to higher levels of yield over the medium to long-term. If there is no reduction in the amount of energy available, the results are strongly influenced by the share of cereals for animal feed and biofuel production. In any case, a change in these assumptions would not change the results in terms of quality, leaving unchanged the ranking of the scenarios in terms of calories produced and the yield.

In the case of an energy crisis, the results greatly depend on the amount of time spent by the systems in shifting from the HEI agricultural model to the LEI model (in the direction of a Stopped HEI Growth scenario). In fact, in the short-term, the results of Strong HEI Growth-Energy Shock and BAU-Energy Shock are less negative.

The example shown here highlights the fragile nature of the global agricultural system. This frailty will have to be tackled positively by promoting a balanced mix of agricultural models, designed to cope with the phenomena of relative scarcity. Of course, the reality is much more complex than what has been willfully represented in this simulation for dissemination purposes. In addition to possible energy shocks, in fact, there are many other long-term risk factors: water availability, adaptation to atmospheric phenomena, etc.

However, in any case, the result presented is by no means trivial because it does illustrate one of the issues that will be very important in the future, when the search for solutions based on approaches to energy consumption content and an in-depth knowledge base (according to the logic of balance already described) will become one of the most crucial aspects of sustainability.





#### The future of agriculture

Common Agricultural Policy (CAP).

ion defined its strategic guidelines and agriculture.) directions of the policy for orienting the With regard to environmental protecnext decade.

In particular, the EU is considering us- greenhouse gas emissions caused by ing the CAP reform to establish a new agriculture, even though there has aldistinctive features of this have been We are also giving greater attention to ward 2020: Meeting the food, natural ing of the many rural areas still highly resources and territorial challenges of dependent on the primary sector for its

and strengthen its production capac- demand for food commodities. text. Secondly, there are two further the close observation of the evolution

In November 2010, the European Un- of rural areas and the diversity of EU

of direct payments to practices which

techniques could contribute.

tives defined in the overall European Strat- territorial challenges of the future. egy 2020, the CAP, in particular, should:

- where agriculture is an important eco- markets in times of instability.

more efficient and more effective."

- mate change;
- 3rd objective: territorial balance.

also the more efficient use of scarce. A more detailed explanation of the main resources and the mitigation of the im- technical options currently at the focus of pacts of climate change to which these the debate can be found in the Commission document: The CAP toward 2020: In light of this analysis, and given the objec- meeting the food, natural resources and

At this point one might ask what obserreached with a view to ensuring the the BCFN has produced and produces on

global demand for food commodities; eral occasions by the BCFN, the issue produced in a sustainable way, in ac- challenge of food security, which is adcordance with the objectives that the dressed through innovation and greater maintain viable rural communities, tutions to intervene in the agricultural

environmental and territorial benefits. environmental and economic dimenand Rural Development - Dacian Ciolog as part of an overall effort to create a the guidelines developed so far when rope. The Commission is proposing €4.5 "European partnership for research and In summary, the three major challeng- innovation." This is aimed at developing es of the new CAP identified so far are: a strategy for sustainable increases in The CAP review affords the EU the oplenging international markets.

crucial aspect. Precisely for this reason, and environmental impact.

be one of the main conduits for encour- recent years. extremes of price volatility.

toward the sustainability of agricultural For that reason, the Barilla Center for models. This should shape the choices Food & Nutrition can express support viding a reference point to orient agri- CAP proposal which suggests safety cultural sustainability. In addition, such a nets (intervention and private storage) process could lead to a greater appre- for the agricultural aspects of the sup-European food production in the chal- Furthermore, transfers could help to spread awareness of the importance of In discussing sustainable agriculture, we respecting, rediscovering and increasmust not ignore the role of biofuel produc- ing the value of local traditions and tion" and "energy target" for the use of the different contexts within the local comscarce terrain, this production could be a munity, be it for tradition, eating habits

particular attention should be given to the The intention of the European Com-In line with the Barilla Center for Food the continued market orientation of the & Nutrition's long-held position, the in- CAP, is positive and seems to be in line tention to revitalize rural communities with the reflections made by the Barof Europe, provides an added value that Last but not least is the need for longpossible to show that operational tools a topic on which the Barilla Center for such as cash transfers to farmers could Food & Nutrition has been very vocal in

practices. In addition, it may assist in uable. This needs to focus on improving one of the challenges of food security velopment of measures to ensure food whic's are of such concern for the future. security and quality, the development It is worth remembering that the Barilla of sustainable agricultural techniques food, starting with those most related

to territorial specificities. EU funds for researchers to farmers, to the inclusion research and innovation and will focus of farmers throughout the process. and farmers.

closer cooperation between scientists of enlargement of the agricultural sector and therefore of the planning of the There are indications of a shift away processors and users that are part of





## 70

# 5. CONCLUDING REFLECTIONS

AGRICULTURE IS A
COMPLEX AREA OF
ACTIVITY THAT DOES NOT
LEND ITSELF TO EASY
SIMPLIFICATIONS

Agriculture is complex and does not lend itself to easy simplification, which is why it is difficult to draw unequivocal conclusions from this analysis. Nevertheless, some basic principles can be identified. This is a collection of evidence, reflections and guidelines that characterize a possible approach for creating true sustainability. In particular, there are seven points of attention that we consider essential and which are stated briefly below:

#### **1** Sustainable agriculture is characterized by a systemic conceptual and operational approach.

For a future of sustainability, we must increasingly learn, according to a multidisciplinary approach, to "hold together" the social, environmental and economic aspects, as well as those of research and development. Approaches aiming at pursuing partial goals can only reap some short-term success of one of the aspects and consequently do not help in winning the challenge of sustainability. An example of this is the intention to revitalize rural communities, an essential condition for effective land management and land use. The richness/diversity of agricultural systems is, in fact, an added value in the perspective of combining efficiency and agricultural sustainability.

#### 2 Sustainable agriculture is based on a large number of already known agricultural practices.

The knowledge available, made up of scientific knowledge and proven practices, has crystallized into some excellent and practical guiding principles of truly sustainable farming. There is, in fact, an ever-growing convergence around the best agricultural practices, specifically declined in different situations, which should be followed. These are 1: systemically use crop rotations (as was standard practice in the past) on the same tracts of land; cultivate a wider range of plant species to arrive at a correct spatial distribution of trees, shrubs, pastures and crops in order to improve the resilience of the system; minimize mechanical working of the ground, in order to maintain the soil structure and organic matter; improve and maintain a protective cover on the surface of the organic soil using species of reduced cycle in the times between crops; and use cover crops or organic crop residues, in order to protect the soil surface, conserve water and nutrients, promote biological activity of the soil and contribute to the integrated management of pests and weeds. These techniques - associated with the use of crop varieties with high yield (resistant to biotic and abiotic stresses and with good nutritional value), the optimized use of organic and inorganic fertilizers, integrated management of pests and diseases through appropriate practices<sup>2</sup> (based on biodiversity, the selection and use of pesticides with low environmental impact) and, when necessary, efficient management of water resources - allow for better performance in terms of sustainability, with the same macro model of reference (HEI, LEI, IEI).3 In Chapter 3 of this paper, there is a brief

summary of the experiment conducted by Barilla through the recovery of these "good" guiding principles at the farms of some of its suppliers of raw materials. At present, the results are very encouraging.

#### **3** Agronomic "knowledge" is not very widespread.

With the development of science, farming is increasingly characterized by the articulation and breadth of knowledge gained regarding the characteristics of the natural environment and the physiology of plant species. All this is combined with the practical experience accumulated over centuries of activity. In other words, there is a wealth of knowledge available of extraordinary value that is only partially used today. In certain surroundings, this seems to be due to the lack of effective processes for transferring knowledge. In others, this is because it is believed that the available technology, at least partially, makes in-depth knowledge of the natural dynamics superfluous. In summary, it could be argued that regardless of the model adopted (HEI, LEI, IEI), the biggest problem that global agriculture now faces is the need to strengthen its human capital base, bridging the gap between available knowledge, individual and system skills. With regard to this aspect, significant investment plans will be needed, because they are the prerequisite for any development in the direction of greater sustainability.

#### **4** Correct agricultural models for specific contexts: the objective is to reduce external inputs.

That said, there are no good or bad agricultural models a priori. There are certainly HEI models that we believe will prove unsustainable and LEI models that cannot be implemented in all contexts; however there is a wide range of realities, namely, of IEI models adjacent to LEI models that can be managed appropriately, in light of the above requirements of sustainability.

The choice of model depends on the context. In geographical contexts in which the HEI systems with high economic performance are rooted (such as the U.S., Brazil and Argentina), it makes no sense to propose or suggest extreme choices of rupture and discontinuity. Instead, the adopted model's limitations on sustainability should be evaluated to make the necessary corrections. Likewise, Europe's way should be implemented with the IEI/ LEI models, based on effective enforcement mechanisms of the application of knowledge. In other words: what matters is the trend line, i.e., the shift toward more sustainable IEI models and the equilibrium with the models within the macro-regions.

A different reasoning must be made for developing countries where there are still no active agricultural models that are economically and socially sustainable; one has to return to and adapt models that are appropriate to the specific characteristics of the local reality.

#### **5** Biodiversity as an instrument for correct risk management.

A pragmatic approach, without prejudice, to the choice of agricultural models allows – at the level of policymaking – the maximization of the overall resilience of agricultural systems. Proper management of biodiversity and the coexistence of different models, all equally optimized concerning sustainability boosts the possibilities of response to adverse events and the search for specific objectives of the system, when these are alternative (e.g., maximum quality vs. large volumes).


#### **6** Investments in technology to make agriculture more able to adapt to change.

According to the reading proposed within technology also takes on a different connotation from that which is too often prevalent in these times. Today, when it comes to technology in agriculture, one is often referring only to productivity and yields, thinking

HIGH PRODUCTIVITY

LOW PRODUCTIVITY

Figure 5.1. The transition toward sustainability



Source: elaborated by The European House-Ambrosetti, 2011.

that these may be increased by improving the individual varieties. However, even more important is the adaptability that is expressed in the integrated and harmonious management of a wide range of tools and logic: plant varieties resistant to stress, management of advanced systems of irrigation, scientific approach to fertilization, etc.

7 The external factors of sustainability in agriculture: food waste, losses, and biofuel.

One cannot forget that a large part of the issues that plague agriculture and the agrofood system lie outside the choice of models and the search to optimize them. As seen
in the opening, there are phenomena of great impact that affect the goals agriculture
has set for itself, emphasizing beyond measure the matter of production volumes, at
the expense of a more balanced approach. There is, above all, the issue of food waste,
which is truly disturbing in its proportions and which is one of the challenges to sustainable agricultural in the future. Along with the matter of loss/waste that world agriculture produces today, there emerges a question that seems central to the choices
regarding the allocation of resources in agriculture (both financially and physical): the
production of biofuels.

For the issue of "waste," and "biofuels," the inadequate management of the problem, and questionable choices in the field of energy policy have resulted in strong pressure on the agricultural system to make up for deficiencies that it should not have to bear.



© Corbis



# HYPOTHESES, ASSUMPTIONS AND INTERMEDIATE RESULTS OF THE BCFN-MILLENNIUM INSTITUTE SIMULATION MODEL

#### Premise

The Barilla Center for Food & Nutrition is conducting a study on the long-term impacts (until 2050) of different agricultural practices (or agricultural models). The study examines different scenarios with the aim of facilitating the understanding of the key mechanisms that link agriculture to the rest of the socio-economic-environmental system and its long-term implications. The initiative is part of a more extensive program of studies relating to the environment and sustainability, and this study on sustainable agriculture will provide a broader systemic perspective and ideas for the upcoming debate on the renewal of the EU's Common Agricultural Policy.

The Millennium Institute (MI), which supports the study of the long-term impacts (until 2050) of different agricultural practices (or agricultural models) conducted by the Barilla Center for Food & Nutrition, for this purpose, has developed and used a Threshold21 (T21) global model for the simulation of scenarios representative of longterm agriculture worldwide (1970-2050), taking into account agricultural development based on the different degrees of diffusion of the model that is currently prevailing (High External Input systems, or HEI) compared to the alternative model (Low External Input, or LEI). Each model is characterized by the amount of resources needed to support it, productivity and the environmental impacts involved. The simulations take extreme conditions into account in terms of resource limitations in the long-term (i.e., energy restriction), in order to assess the adequacy of the different models under these conditions.

The analysis particularly focuses on the aspect of teaching: the goal is not to provide exact predictions or highly probable scenarios, but rather, to examine plausible extreme situations and provide information on key systemic mechanisms that play a fundamental role in shaping the development of agriculture in the long-term.

In summary, the study has the following purposes:

- Evaluate, through an analysis of the scenarios, the implications for the long-term sustainability of the two extreme agricultural models (HEI vs. LEI).
- Highlight how agriculture falls within the broader socio-economic-environmental system.

This report provides a synthetic view of the structure of the model and the general hypotheses illustrates the fundamental assumptions that characterize the different scenarios, and describes and discusses the results of the different scenarios. The work is aimed at a wide audience, which is why this report was written in such a way as to avoid the technicalities of the modeling as much as possible.

### Model Structure and Assumptions

#### General structure and linkage between sectors

This study is global in nature and the world will be treated as a single entity without further geographical breakdowns: the time horizon between 1970 and 2050. Therefore, the modeling focuses particularly on the mechanisms that regulate long-term, socio-economic-environmental development in general, with a specific interest in the agricultural sector. Accordingly, the description of this sector will be more detailed than that of the

The model is composed of 10 sectors, all interacting dynamically, that provide an integrated perspective on long-term development. Figure A.1. illustrates the general structure of the model and the links between the sectors. For simplicity, only the main links are represented. More details are given in the following section.

**EMPLOYEMENT POVERTY** PRODUCTION: AGRICULTURE, INDUSTRY, SERVICES POPULATION LAND HEALTH EDUCATION

Figure A.1. Overview of the model structure: sectors and interactions

Source: Millennium Institute, 2011.

The 10 sectors have the following key features:

- Population: divided according to age (81 classes) and sex. The fertility rate is determined by income (production sector) and education (education sector), while the mortality rate is determined based on life expectancy (health sector).
- Education: measured in terms of average years of schooling and determined on the basis of the total expenditure on education per pupil, which in turn is defined in terms of the share of the global GDP.
- Health: measured in terms of life expectancy at birth, separately for men and women. Life expectancy is determined by income (production sector) and the overall health expenditure per capita, in turn calculated as a share of the global GDP.
- Employment: represented separately for agriculture, industry and services sectors. For industry and services sectors, employment levels are calculated according to the level of physical capital (production sector) and education (education sector - higher levels of

education imply greater capital intensity). For agriculture, employment is also determined by the land cultivated (land sector).

- *Poverty*: measured in monetary terms (percentage of population below the poverty line) and in terms of nutrition. With regard to nutrition, the average levels per capita of calorie, protein and fat intake are determined on the basis of agricultural production (production sector). Moreover, the percentage of the population below the minimum levels of energy intake in the diet is determined using a log-normal distribution approach.
- Production: includes agricultural production, industrial production and services; it is determined by using an extended Cobb-Douglas production function with endogenous determination of the total productivity of the factors (PTF). For industry and service sectors, the key inputs are labor (industry employment) and capital; productivity is determined by the level of education (education sector) and health (health sector). For agriculture, the production is divided further into different activities and crops, as explained in more detail in the Appendix.
- Land: this sector considers 4 types of land areas: agricultural area, forest area, settlement area
  and other land (which includes all other land types not included in the previous aggregates).
   The growth of the agricultural and settlement areas is determined by population growth (population sector) and is limited by the small proportion of forest land and other land converted for such purposes. According to the FAO classification, the agricultural area is divided further into arable land and permanent crops, as well as permanent pastures and prairies.
- Water: this sector determines the use of water for agricultural, industrial and domestic/municipal activities. The demand of water for agricultural use is calculated based on cultivated area (land area) and the type of crop (production sector). The demand of water for industrial use is determined by the level of industrial production (production sector). The demand of water for domestic/municipal use is determined by the total population (population sector) and per capita income (production sector).
- Energy: the energy demand is divided into 5 types: oil, gas, coal, electricity from non-renewable sources and electricity from renewable sources. The demand for energy is broken down into the following sectors: agriculture, industry, services, transportation, residential transportation and others. The energy demand is based on the intensity of capital (production sector), the total population (population sector), income per capita (production sector), as well as on energy prices, the extent of network coverage and technical progress, defined as exogenous factors.
- *Investment*: the economic resources that are saved are invested in different sectors according to their relative size and the relative profitability of each sector.

#### Structure of Agriculture

Given the primary purpose of the study, an analysis of scenarios in relation to alternative agricultural models -, the agriculture sector has been described with a particular wealth of detail. Similarly, the sector on poverty concerning nutrition and the distribution of food is more in-depth than other sectors. The following paragraphs provide a description of the specific components of the model. Agricultural production is divided into crop production, production of food of animal origin and forestry production. The production of crops and foods of animal origin is subdivided further as shown in Figure A.2.

The 11 categories related to the production of crops cover the full range of global agricultural production, while the last remaining category – Other – represents only a very limited part of the production and mainly includes products that are immaterial in terms of nutrients (such as spices).

Figure A.2. Suddivisions of the production of crops and foods of animal origin

CROP PRODUCTION	PRODUCTION OF FOOD OF ANIMAL ORIGIN
Cereals for food (including rice, wheat and millet)	Meat
Cereals for fodder (all the other cereals)	Dairy products
Fibrous crops	Eggs
Fruit	Fish (exogenous)
Oilseed crops	
Legumes	
Roots/tubers	
Nuts	
Vegetables	
Sugar crops	
Other	


Source: Millennium Institute elaboration of FAO data, 2011.

The production is determined for each category of crops according to 9 main factors (Figure A.3.): physical capital, energy, soil quality, water, cultivated area, knowledge, education, health and employment. While most of these factors are determined in the areas described above, the availability of knowledge in the agriculture sector and of soil quality is calculated in specially processed, specific sub-sectors.

The combination of knowledge about farming techniques and seed varieties and fertilizers is considered a key component of knowledge in the agricultural sector. The accumulated expenditure for R&D in agriculture is used as a proxy for such a combination of knowledge. Private and public R&D investments, determined in terms of shares of added value generated by agriculture, are listed separately. With regard to soil quality, the density of macronutrients in the top layer of soil is used as a reference, specifically nitrogen, phosphorus and potassium. The nutrient density decreases with the growth of crops and increases with the use of inorganic and organic fertilizers and other organic methods.

The production of meat, dairy products and eggs is determined according to the area designated for permanent pastures and meadows, the quantity of feed grain produced,

Figure A.3. Key resources that influence agricultural production




Source: Millennium Institute, 2011

and the productivity of agriculture. The production of fish is exogenous, as it does not fall within the scope of this study.

The flow of the production of food crops (thus excluding fibrous crops) and foods of animal origin are used to determine the average levels of nutrition, including the amount of calories, proteins and fats per person per day. To calculate these levels, the agricultural productions - net loss of crop -, other uses and losses are further subtracted (Figure A.4.): losses during processing, the amount used for sowing (if applicable), the amount used for animal fodder (if applicable), the amount used to produce biofuels and other non-food products and the losses incurred during shipping and distribution. What remains is the amount of agricultural production that is actually available for human consumption. Therefore, the average levels of nutrition per capita are determined by applying the corresponding average content of nutrients in each of the 10 crops and each of the four food products of animal origin.

The average levels of nutrition are also used to determine the distribution of food by applying a lognormal approach. This allows one to calculate the proportion of the population below the minimum dietary standards. In addition, it determines the proportion of food waste at the household level in order to calculate the overall proportion of food that is not consumed.

Figure A.4. Flow of agricultural food production in 2009



Source: FAO data base

#### Agricultural models and scenarios


#### Characteristics of different agricultural models

The FAO (2011) provides a first, broad definition for the two types of agricultural models considered in this analysis, supplemented further by additional assumptions described herein. The FAO defines the HEI and LEI agricultural models as follows:

- High External Input (HEI) systems:
- Orientation to the commercial market;
- Use of high-yielding improved varieties;
- Mechanization with low human labor intensity;
- Almost complete reliance on external inputs of synthetic nature (fertilizers, pharmaceuticals, etc.).
- Low External Input (LEI) systems:
- Broader trend toward subsistence and less oriented toward the market;
- Use of traditional cultivars;
- Techniques with high content of knowledge and labor;
- Little or no use of external nutrients, no use of synthetic chemicals to control pests and diseases, but strong emphasis on nutrient cycles on-site.

In fact, there is a continuum of combinations of HEI and LEI practices and, therefore, it is not possible to classify a real case as belonging entirely to the HEI or the LEI system.

Figure A.5. Overview of different LEI, IEI and HEI system, according to the FAO



Source: FAO/OECD, Food Availability and Natural Resource use in a green economy context, 2011.

As already noted above, the FAO identifies a wide variety of farming systems with respect to the relative intensity of use of external inputs.

For simplicity, the analysis quantitatively characterizes the two different agricultural models (LEI and HEI) as follows:

Hypothesis 1: the Low External Input (LEI) farming system uses about 35% more labor per hectare of cultivated land than the High External Input (HEI) farming system.

This hypothesis is based on the results of the following studies:

- The Soil Association (2006) calculated that organic agriculture in the United Kingdom provided 32% more jobs per farm than conventional farming.
- In Denmark, a conversion from conventional agriculture to organic agriculture increased the need for human labor by 35% (Barthelemy, 1999).
- A study carried out in Turkey on the production of raisins in 82 conventional and organic farms concluded that the inputs of human labor were on average higher (approximately 10%) for the organic farms (Gündogmus et al., 2006).
- Organic systems need about 15% more labor on average (Sorby, 2002; Granatstein, 2003).
- In addition to performing tasks of weeding, cultivation and care of plants and animals (largely done with machinery and chemicals in conventional systems), farmers who rely on the organic system plant cover crops, spread manure and produce fertilizer (FAO 2007).

Hypothesis 2: farming with the Low External Input (LEI) system uses about 50% less energy per hectare of cultivated land than farming with the High External Input (HEI) system.



- According to the Soil Association (2006), the largest share of energy used in conventional agriculture on average, 37% of the total was represented by synthetic pesticides and mineral fertilizers, especially nitrogen and, to a lesser extent, phosphorus and potassium.
- Refsgaard and his collaborators (1998) found that the consumption of energy associated with the use of fertilizers represented between 25% and 68% of total energy use, depending on the type of crop and growing conditions.
- A study carried out in Turkey on the production of raisins in 82 conventional and organic farms concluded that the average energy inputs were lower (around 38%) on the organic farms (Gündogmus *et al.*, 2006).
- Pimentel (2005) estimated that between 28% and 32% less energy is used in organic farming.
- Farming practices and the use of machinery greatly affect the use of energy on individual farms, but there is no evidence that organic farming requires less energy for the mechanical processes. For example, several studies reported that the production of organic potatoes and carrots requires high energy input per unit of output due to mechanical weed control (Stolze *et al.*, 2000; Williams *et al.*, 2006; Bos *et al.*, 2007).
- Typically, organic agriculture production uses 30 to 50% less energy than comparable non-organic farming (FAO, 2007). Although organic farming uses energy more efficiently on average, it often requires an indirect trade-off of energy-intensive inputs with additional hours of human labor, about a third more than conventional farming.

*Hypothesis* 3: the density of nitrogen in the top layer of soil in the Low External Input (LEI) agriculture system is about 30% less than in the High External Input (HEI) agriculture system and, therefore, the yields of the LEI farming are lower.

- The crop rotations with legumes in the LEI system resulted in an intake of nitrogen in the model, derived mainly from biological nitrogen fixation of 90 kg per hectare per year. This contribution was equal to only 35 to 50% of that for a crop in the HEI system and, therefore, the yields were correspondingly lower (Wolf, 2002).
- Comparison of productivity of LEI vs. HEI: less than about 9% for the LEI system (Stan-hill 1990)
- Comparison of Productivity of LEI vs. HEI: more than 80% for the LEI system in developing countries and less than about 8% for the LEI system in industrialized countries (Badgley *et al.*, 2007).
- The transition from the HEI to the LEI system is not easy: it gets worse before getting better results (Badgley *et al.*, 2007) and the transition period lasts five years (Pimentel, 2005).

#### The scenarios formulated

The scenarios have been analyzed taking into account agricultural development based on different degrees of diffusion of the currently prevailing model (High External Input, or HEI systems) compared to the alternative model (Low External Input, or LEI systems). Each model is characterized by the amount of resources needed to support it, the productivity, and the environmental impacts involved. The simulations have taken into account the extreme conditions in terms of resource limitations in the long term (i.e., limitations in terms of energy resources, water and land) to assess the adequacy of the different models under these conditions.

There is the intensive use of inorganic fertilizers as a proxy to estimate the prevalence of past and present HEI systems. Based on that estimate, in 1970 approximately 30% of the total cultivated area was managed with HEI systems, and this value reached 45% in 2010. Based on the current trend, it is possible to formulate several scenarios, including:

- Scenario of Business As Usual (BAU): the practices with a high level of external input continue to spread as in the past, coming to cover 60% of the total cultivated area in 2050;

e

- Scenario of Strong HEI Growth: the practices with a high level of external input spread at an accelerated pace to cover 90% of the total cultivated area in 2050;
- Scenario of Stopped HEI Growth: practices at a high level of external input do not spread any further and their coverage remains constant at the current level (45% of the total cultivated area).

These three scenarios are examined in the light of two different sets of assumptions in relation to the availability of energy, and more specifically:

- all the normal scenarios (I): oil prices rise gradually to reach \$130/ barrel in 2035 (IEA average projection), then grow further to reach \$170/barrel in 2050 (dollar figures in real terms of the year 2001);
- all the normal scenarios (II): inorganic fertilizer prices remain affordable for most farmers, thereby not inhibiting their spreading;
- all the Energy scenarios (I): oil prices rise faster than the reference value between 2025 and 2030, reaching \$200/barrel in 2030, then increasing further to reach \$280/barrel in 2050 (dollar figures in real terms, 2001);
- all the Energy scenarios (II): due to the rapid increase in oil prices, the prices of inorganic fertilizers increase substantially with a consequent reduction in their use: no more than 50% of the total cultivated area is fertilized in areas where the farms have a higher added-value potential.

Finally, all scenarios put forward the following hypotheses regarding the use of agricultural production:

- A gradual increase in meat production per capita: from 40 kg/person/year in 2010 to about 65 kg/person/year in 2050 (BAU);
- An increase in the share of cereals (except rice, wheat and millet), oilseed crops and sugar crops used for biofuels and other non-food purposes:
- Cereals: from the current 6% to 10% in 2050;
- Oilseed crops: from the current 7.5% to 15% in 2050;
- Sugar crops: from the current 0.5% to 2% in 2050.

#### Scenario Results

#### General socio-economic and environmental developments in the BAU scenario


The following is an overview of the results of the Business As Usual scenario (BAU) in the 1970-2050 period. For the 1970-2010 period, series of historic data on over 250 indicators were collected, with the aim of comparing them with the results of the model. Therefore, the results of a series of general indicators are discussed, which include: total population, life expectancy, average duration of schooling, the real rate of the GDP growth, GDP per capita, agricultural area, forest area, total water demand and overall energy demand. The time-line graphs show multiple lines that include a series of data and model projections. In the presence of multiple indicators, additional lines are highlighted with different colors (see figure legends). In some cases, when there are authoritative projections up to 2050, the line of historical data also extends up to 2050 (from 2010 to 2050).

However, the simulation results should not be construed as forecasts, but rather, as a projection obtained from a series of basic assumptions.

In summary, the BAU scenario indicates that by 2050 the global population will slightly exceed 9 billion people, a slightly higher figure than the one calculated by the UN Population Division (Figure A.6.). This finding is consistent with the pattern of life expectancy at birth, which gave slightly higher projections (about 80 years for women in 2050). In addi-

tion, there is a substantial increase in the average duration of schooling, which in 2050 will come to be roughly 7.25 years.


Figure A.6. Population and life expectancy from birth



Source: data/projections by UN POP and T21 projections.

While these social developments are positive, they are, at the same time, both the result and the cause of rapid economic growth. With regard to the GDP per capita, real growth is estimated to be 2.7% in 2030, before falling gradually back to about 1.25% in 2050 (Figure A.7.). This rapid increase is supported, in particular, by the growth in the industry and services sectors, which appears rather slow (in terms of added value) in the agricultural sector.

Figure A.7. Real growth of the GDP per capita



Source: World Bank data and T21 projections


In terms of land use, the BAU scenario foresees a stabilization in the use of land for agricultural purposes, which in turn implies a consolidation of the reserves of forest areas. However, stabilization in the dynamics of the land does not slow the growth of the agricultural sector, as shown later. It is expected that, in the long-term, the demand for water and energy will increase substantially, reaching 6,300 km<sup>3</sup>/year and 750 quadrillion BTU/year, respectively, in 2050.

#### BAU Results for the Indicators Related to Agriculture

Here we discuss the results for specific indicators of agriculture, including: the yield of cereals, calories per person per day, the percentage of the population that is undernourished, water demand for agriculture, and energy demand for agriculture.

According to projections, the yield of cereals for food and those for fodder will have similar rowth rates, driven by increasing mechanization, higher human capital, more in-depth knowledge in the sector and a higher concentration of nutrients in the surface layer of the soil. More specifically, it is expected that in 2050 the yield of cereals for food will rise to 5.6 tons/ha (from the current 3.3), while that for the fodder sector will reach 6.2 tons/ha (from the current 3.7) (Figure A.8.).

Figure A.8. Yield of cereals for the food industry and for the fodder industry




Source: FAO data and T21 projections.

The combination of the increase observed in yield and a gradual slowdown in population growth leads to a continuous increase in the amount of calories available on a per capita basis (Figure A.9.).

The projection indicates approximately 3145 kcal/person/day of foods for human nutrition by 2050. So sudden an increase of calories available per capita leads to a smaller percentage of the malrnourished population, which, by 2050 will have dropped to around 3.5%. Translated into absolute figures, by 2050 the number of individuals who suffer from hunger will have decreased to 345 million. This BAU projection depicts a rapidly changing agricultural sector, which changes to meet the needs of the growing world population and will eventually be able to generate almost enough output to overcome hunger in the world.

In contrast, the BAU scenario also implies a greater impact of agriculture on the natural environment. In particular, it is expected that the demand for energy in agriculture will grow from the current 7 quadrillion BTU/year to about 8.5 in 2050. A similar increase in the demand for water is also expected (although at a more moderate speed), so that the current value of 2.8 km³/year will grow to about 3 in 2050. The more modest increase of the water demand is due to the stabilization of the cultivated areas, the gradual increase in the percentage of irrigated areas, and the continuous improvement of water management.

Figure A.9. Production of calories per capita and percentage of malnourished population



Source: FAO data and T21 projections.

#### Comparative Analysis of the Results of different Scenarios

The following presents the results of the alternative scenarios (BAU, Stopped HEI Growth, Strong HEI Growth), first assuming the absence of a future limitation of energy and then, assuming a substantial limit linked to the availability of energy.

We now analyze the results of a specific indicator that summarizes the issues related to agricultural production and the use of the product: calories per capita from foods for human nutrition. To simplify matters, the results of a single indicator are shown (although the model gives rise to several important indicators of nutrition) and, specifically, the calories of food per capita have been chosen because of their importance in human nutrition.


As mentioned above, assuming that there are significant restrictions related to the availability of energy, the BAU scenario shows a steady growth in the amount of calories produced for human nutrition, which in 2050 will reach 3145 kcal/person/day (Figure A.10.). This data is in line with the FAO projections for 2050, equal to 3130 kcal/person/day (FAO, 2011).

In the Stopped HEI Growth scenario, productivity growth is slower as compared to the BAU scenario, because inorganic fertilizers are used in smaller quantities. As a result, the increase in calories produced for human nutrition will be lower, and by 2050, will have reached 3015 kcal/person/day.

Due to the higher concentration of macronutrients being artificially added to the soil, in the Strong HEI Growth scenario the amount of calories produced for human nutrition is growing faster and will have reached 3410 kcal/person/day in 2050.

In the event of a substantial decrease in energy availability starting in 2025 (all the Energy

Figure A.10. Agricultural production for human nutrition



Source: FAO data base

Shock scenarios, Figure A.11.), the increase in the amount of calories for human nutrition would continue unchanged until 2025, but would then slowdown in different ways according to the specific scenarios.

In the BAU-Energy Shock scenario, there is a gradual decrease in the amount of calories produced for human nutrition beginning in 2025 before stabilizing and then seeing growth again toward 2035.

This decrease is due to the combination of two factors: first, the increase in oil prices that has direct negative effects on productivity and yield, because as the cost increases, the use of oilpowered vehicles (such as tractors, water pumps, etc.) decreases; and secondly, the reduced access to fertilizers forces some farmers, who in other circumstances would practice a HEI type of agriculture, to switch to LEI (in the direction of a Stopped HEI Growth scenario) farming. This forced switch to the LEI system is an expensive process because part of the existing physical capital must be substituted/adapted and because it requires the acquisition of additional knowledge.

In addition, conversion to LEI farming is an expensive process in terms of time: it inevitably takes some time before the farmers realize that the increase observed in energy prices is not transitory, and before they acquire the human and physical capital needed to implement the LEI practices. These resources might not be immediately available at the time of rising energy prices and declining productivity.


At the same time, when the LEI production systems have not yet been implemented, but the fertilizers are still available in the quantities needed, the land is further impoverished.

Finally, once the LEI practices are implemented, it takes about five years before the soil is fertile again. Overall, it is assumed that the average transition from HEI to LEI farming requires about 10 years, given the conditions of limited access to energy resources and in-

organic fertilizers. This period represents the loss of a decade in terms of increased yields. The forced transition from HEI to LEI has more drastic effects in the Strong HEI Growth-Energy Shock scenario; in this case the yield and, consequently, the amount of calories produced for human nutrition suffer a substantial decline in 2025, before resuming growth in the last 15 years of the scenario. The most significant decrease in yield is due to the fact that in this scenario, a higher percentage of farms must change from the HEI to the LEI one, since for the HEI scenario we assume there is a higher final level of dissemination of the HEI practices (90% in 2050).

Finally, the impact of the sudden rise in oil prices and reduced access to inorganic fertilizers is lower than in the Stopped HEI Growth-Energy Shock scenario. This is due to the fact that in the Stopped HEI Growth scenario, a smaller percentage of the cultivated area is managed according to HEI practices: the level estimated for 2050 remains at the current 45%. In this context, the 50% reduction in the availability of inorganic fertilizers does not represent a limitation for agricultural productivity because the target level of distribution of HEI practices remains below 50%. Therefore, the decline in the growth of yields that is observed is only due to the direct impact of the increase in oil prices on the cost/use of mechanical means powered by oil. The simulation results showed that, if after 2025 substantial limitations of energy availability should be detected, an approach of low external input could lead to a Worse-Before-Better result (WBB), with lower productivity in the short-term but increased productivity in the long-term. The results of the simulation of the scenarios without substantial energy limitations are significantly influenced by the assumptions regarding the amount of cereals used for fodder and the amount of crops grown for biofuels. However, any changes in these assumptions do not entail a substantial change in the results in qualitative terms: the ranking of the scenarios in terms of yield and calories produced remains unchanged.

Figure A.11. Agricultural production for human nutrition in case of an energy shock



Source: FAO data base

The results of the simulation of the Energy Shock scenarios are particularly sensitive to the amount of time assumed to be necessary for the transition from HEI to LEI. In fact the results of Strong HEI Growth-Energy Shock and BAU-Energy Shock scenarios are less negative. However, even if the transit time were significantly lower (five years), the positioning of the scenarios in terms of yield and calories produced would remain the same (the difference in quantitative terms is still quite low).

#### Summary and Conclusions of simulation

In summary, in the BAU scenario, agricultural production grows at a constant rate, generating a continuous increase in the amount of calories per capita for human nutrition. This leads to a significant reduction of the population that is malrnourished, reaching 345 million in 2050 (3.5%). The HEI scenario generates an even higher level of yield and, consequently, increased production for all crops; while on the contrary, the Stopped HEI Growth scenario generates yields and production that are slightly lower than with the BAU system. This is mainly due to the lower density of nutrients in the soil that characterizes the LEI type of agriculture as compared to that of HEI. Consequently, the amount of food calories available per person per day is higher in the HEI scenario compared to the BAU system, whereas it is lower in the LEI scenario. The relative difference between the scenarios is less when considering the quantity of the population that is malnourished, which means that the additional production generated within the HEI scenario is only partly for the benefit of the poor.

On the contrary, the Strong HEI Growth scenario has a greater environmental impact than the BAU scenario, particularly with regard to energy demand. In the Energy Shock scenario, it is hypothesized that the more intense exploitation of the natural resources would not be practical due to the limited availability of those very resources. What follows is a rapid reduction of the yields for the Strong HEI Growth-Energy Shock scenario, with lower levels than those of all the other scenarios. Therefore, agricultural production grows slower in the Strong HEI Growth-Energy Shock scenario compared to the BAU-Energy Shock and Stopped HEI Growth-Energy Shock scenarios, and the quantity of calories available per person by 2050 is less compared to the other scenarios.

This figure can be traced to a number of factors. First, farmers do not react immediately to changes in prices, but expect to see if the changes are temporary. Secondly, to change farming practices it is necessary to acquire a certain amount of knowledge on LEI agriculture. Thirdly, the available capital may not be suitable for LEI agriculture and may need to be replaced. And finally, the transition from monoculture to polyculture involves changes in marketing structures that, in turn, also require time. It is hypothesized that this adaptation results in an average delay of 10 years, with a faster reaction time in countries of medium/ high incomes and longer reaction times in low-income countries.

The consequence of this delay in adapting is that the HEI practices are continued for several years under unfavorable conditions (for example, in the absence of fertilizers), with a consequent impoverishment of the soil. Under these conditions the capital accumulated so far in terms of R&D is, therefore, only partially useful, and thus, productivity drops further. Once the transition from a HEI system of agriculture to a LEI has occurred, the soil has a below-average productivity, and it will take a few years before it returns to being fully productive. In the meantime, a considerable proportion of the potential agricultural production is lost, namely a decade of potential increase in the yield.

The impact of these dynamics on nutrition is important both in terms of the average amount of calories per capita and in terms of the percentage of the population that is malnourished. Just as important is the impact of this transition on the natural resources: energy demand is

substantially reduced. However, this decline is gradual, confirming the fact that the process of replacing and upgrading physical capital is slow and progressive.



© Corbis



# 9!

### **NOTES AND REFERENCES**

#### **EXECUTIVE SUMMARY**

1. FAO/OECD "Expert Meeting on Greening the Economy with Agriculture," Paris, September 5-7, 2011. In precedence, see also IIASA/FAO, "Global Agro-Ecological Zone Assessment input levels," 2010.

#### CHAPTER 1

- 1. Referred to as BCFN.
- 2. "Food production that makes the best use of nature's goods and services while not damaging these assets" (Pretty, 2005).
- 3. FAO. 2008.
- 4. FAO/OECD, "Expert Meeting on Greening the Economy with Agriculture," Paris, 5–7 September, 2011.
- 5. In this regard, see Chapter 4.
- 6. For a comprehensive overview of the major global challenges facing the agricultural sector specifically with the goal of sustainability refer to: *Final Report and Background documents* by the FAO, "International Technical Conference on Agricultural biotechnologies in Developing Countries: Options and opportunities in Crops, Forestry, Livestock, Fisheries and Agro-industry to face the Challenges of Food Insecurity and Climate Change," 2010. For a further discussion of the role of food biotechnology in promoting agricultural sustainability, refer to the position paper *Beyond GMOs. The agro-food biotechnologies*, created by the Barilla Center for Food & Nutrition in 2011.
- 7. World Food Programme, Annual Report 2007.

#### CHAPTER 2

- **1.** FAO/OECD, "Expert Meeting on Greening the Economy with Agriculture," Paris, September 5- 7, 2011. Also see, by the IIASA-FAO, "Global Agro-Ecological Zone Assessment Input Levels" (2010).
- **2.** In Europe, and other countries, regulations exist that codify the distinctive features of organic farming in a clear and comprehensive way.

#### CHAPTER 3

**1.** Gran, Y., C. Liang, X. Wang, B. Mc Conkey, *Lowering Carbon Footprintof Durum Wheat by Diversifying Cropping Systems*, in "Field Crop Research", 122 (3), pp. 199-206, Elseviez, 2011.

#### CHAPTER 4

- 1. Millennium Institute (MI) "is an independent and non-partisan nonprofit organization committed to promoting systems literacy and dynamic modeling tools to attain sustainable development worldwide" (website of the Millennium Institute, "Who We Are").
- 2. In the following pages is a summary of assumptions and results of the model: a detailed description of the methodology used for its construction, hypotheses and underlying assumptions, and the overall results may be found in the Appendix of this document.
- 3. The model of low external input agricultural production, LEI (Low External Input), uses about 35% more work per hectare of land cultivated than a model with high external input, HEI (High External Input). Low external input agriculture (LEI) uses about 50% less energy per hectare than a high-input model (HEI).
- 4. The difference in yield between HEI and LEI is a topic that is still widely debated. Although numerous studies indicate that HEI generally has relatively better yields (Badgley *et al.*, 2007, Stanhill, 1990), the yield of each type of production model depends on the economic, social and environmental context in which it is applied. For the purposes of this study, we introduce the hypothesis that in low external input agriculture (LEI), the density of nitrogen in soils is about 30% lower than in high external input agriculture (HEI); consequently, the yield per hectare is lower in the LEI model. This gap tends to shrink in the long run due to a general improvement of knowledge for the efficient application of the LEI model.
- **5.** Figure 4.1. shows how the amount of calories produced is constantly growing.
- **6.** For further information, see the FAO/OECD study, *Food Availability and Natural Resource Use in the Green Economy Context*, p. 19.

#### CHAPTER 5

- **1**. Save and Grow A Policymaker's Guide to the Sustainable Intensification of Smallholder Crop Production, FAO, 2011.
- 2. IPM, or Integrated Pest Management.
- 3. HEI High External Input; LEI Low External Input; iEI Intermediate External Input.

www.barillacfn.com



